

Some simplicial complexes in combinatorics

[Corrected slides of a talk at UConn Algebra Seminar,
2021-03-11]

Darij Grinberg

August 8, 2025

Abstract. A number of combinatorial identities are concerned with certain classes of subsets of a finite set (e.g., matchings of a graph); they can be viewed as saying (roughly speaking) that equal numbers of these subsets have even size and odd size. In this talk, I will discuss a few such identities – some of them new – and their topological meaning. As a common theme, the "parity bias" (or lack thereof) is the Euler characteristic of a simplicial complex, and thus any expression for it is potentially the tip of a topological iceberg. Underneath are questions of homology, homotopy or even discrete Morse theory. Aside from the specific complexes in question, I hope to provide one more pair of "simplex glasses" through which combinatorial identities appear in a new light.

Preprint:

<https://arxiv.org/abs/2009.11527>

1. Introduction

1.1. Alternating sums

- Enumerative combinatorics is full of *alternating sums*. Some examples:

$$\begin{aligned} \sum_{k=0}^n (-1)^k \binom{n}{k} &= 0 \quad \text{for integers } n > 0; \\ \sum_{k=0}^m (-1)^k \binom{n}{k} &= (-1)^m \binom{n-1}{m} \quad \text{for } m \geq 0; \\ \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} &= (1 \text{ or } 0 \text{ or } -1); \\ \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{ak+b}{c} &= 0 \quad \text{for } c, n \in \mathbb{N} \text{ with } c < n; \\ \sum_{i=0}^m (-1)^i \text{sur}(m, i) &= (-1)^m, \end{aligned}$$

where $\text{sur}(m, i) = (\# \text{ of surjections from } \{1, 2, \dots, m\} \text{ to } \{1, 2, \dots, i\})$.

- These alternating sums are among the most helpful tools in proving identities. (They often play a similar role as the formula $1 + \zeta + \zeta^2 + \dots + \zeta^{n-1} = 0$ for ζ being a nontrivial n -th root of unity plays in the discrete Fourier transform.)
- An alternating sum identity generally looks like this:

$$\sum_{(\text{some finite set})} (-1)^{(\text{something})} (\text{something}) = (\text{something typically simpler}).$$

- In this talk, I shall
 - present some alternating sum identities and their combinatorial proofs by “**toggl**ing” or “sign-reversing involutions”;
 - discuss how a few of these identities can be lifted to **topological** statements about simplicial complexes,
 - and how these topological statements can be lifted to **combinatorial** statements again using discrete Morse theory.
- This is not a theory talk; you’ll hear my personal favorites, not the most general or most important results.
- There will be various open questions.

2. Toggling

2.1. All subsets

- We start with the first identity listed above:

Theorem. Let n be a positive integer. Then,

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0.$$

- There are many ways to prove this:
e.g., expand $(1 - 1)^n$ using the binomial theorem.
- Here is a **combinatorial proof**:
Set $[n] = \{1, 2, \dots, n\}$. Then,

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = \sum_{I \subseteq [n]} (-1)^{|I|}.$$

Claim: In the sum on the RHS, all the addends cancel out.

Proof. For each subset I of $[n]$, we can

1. insert 1 into I if $1 \notin I$, or
2. remove 1 from I if $1 \in I$.

This gives us a new subset of $[n]$, which we denote by $I \triangle \{1\}$.

Easy to see: The map

$$\begin{aligned} \{\text{subsets of } [n]\} &\rightarrow \{\text{subsets of } [n]\}, \\ I &\mapsto I \triangle \{1\} \end{aligned}$$

is an involution (i.e., applying it twice gives the identity), and it flips the sign (meaning $(-1)^{|I \triangle \{1\}|} = -(-1)^{|I|}$ for any subset I of $[n]$).

Hence, all addends in the sum $\sum_{I \subseteq [n]} (-1)^{|I|}$ cancel out (the I -addend cancelling the $I \triangle \{1\}$ -addend). Thus, the sum is 0, qed.

- Our notation $I \triangle \{1\}$ is a particular case of the notation

$$\begin{aligned} I \triangle J &= (I \cup J) \setminus (I \cap J) \\ &= (I \setminus J) \cup (J \setminus I) \\ &= \{\text{all elements that belong to exactly one of } I \text{ and } J\} \end{aligned}$$

for any two sets I and J .

- If a is any element, then the operation of replacing a set I by $I \triangle \{a\}$ (that is, inserting a into I if $a \notin I$, and removing a from I otherwise) is called *toggling* a in I . This is always an involution: $(I \triangle \{a\}) \triangle \{a\} = I$ for any I and a .
- **Remark:** It was actually sufficient for our proof that the map $I \mapsto I \triangle \{1\}$ is a bijection, not necessarily an involution. But all such maps we will encounter are involutions.

2.2. All subsets not too large

- Let us try the second identity:

Theorem. Let n be any number (e.g., a real), and let m be a nonnegative integer. Then,

$$\sum_{k=0}^m (-1)^k \binom{n}{k} = (-1)^m \binom{n-1}{m}.$$

- **Proof.** First of all, we are proving a polynomial identity in n , so we WLOG assume that n is a positive integer (since two polynomials over a field are equal if they agree on sufficiently many points).

We have

$$\sum_{k=0}^m (-1)^k \binom{n}{k} = \sum_{\substack{I \subseteq [n]; \\ |I| \leq m}} (-1)^{|I|}.$$

Now, we try the involution from the previous proof:

$$\begin{aligned} \{\text{subsets of } [n]\} &\rightarrow \{\text{subsets of } [n]\}, \\ I &\mapsto I \triangle \{1\}. \end{aligned}$$

Unfortunately, applying it to a set I might break the $|I| \leq m$ restriction. But it restricts to an involution

$$\begin{aligned} \mathcal{A} &\rightarrow \mathcal{A}, \\ I &\mapsto I \triangle \{1\}, \end{aligned}$$

where

$$\mathcal{A} = \{\text{subsets } I \text{ of } [n] \text{ with } |I \setminus \{1\}| < m\}.$$

Thus, all addends in the sum $\sum_{\substack{I \subseteq [n]; \\ |I| \leq m}} (-1)^{|I|}$ cancel except for those with

$|I \setminus \{1\}| = m$. We get

$$\sum_{\substack{I \subseteq [n]; \\ |I| \leq m}} (-1)^{|I|} = \sum_{\substack{I \subseteq [n]; \\ |I| \leq m; \\ |I \setminus \{1\}| = m}} (-1)^{|I|} = \sum_{\substack{I \subseteq [n]; \\ 1 \notin I; \\ |I| = m}} (-1)^{|I|} = (-1)^m \binom{n-1}{m},$$

since there are exactly $\binom{n-1}{m}$ many subsets I of $[n]$ satisfying $1 \notin I$ and $|I| = m$. This completes our proof.

2.3. Lacunar subsets

- Now to the third identity:

Theorem. Let n be a nonnegative integer. Then,

$$\sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} = \begin{cases} 1, & \text{if } n \% 6 \in \{0, 1\}; \\ 0, & \text{if } n \% 6 \in \{2, 5\}; \\ -1, & \text{if } n \% 6 \in \{3, 4\}, \end{cases}$$

where $n \% 6$ means the remainder of n divided by 6.

- To prove this combinatorially, we need to find out what $\binom{n-k}{k}$ counts.
- **Convention.** We shall write $[m]$ for $\{1, 2, \dots, m\}$ whenever $m \in \mathbb{Z}$.
- **Definition.** A set I of integers is said to be *lacunar* if it contains no two consecutive integers (i.e., there is no $i \in I$ such that $i+1 \in I$).
- For example, $\{1, 3, 6\}$ is lacunar, but $\{1, 3, 4\}$ is not. Empty and one-element sets are always lacunar.
- Note that any lacunar subset of $[n-1]$ has size $\leq \lfloor n/2 \rfloor$.
- **Proposition.** For any $n \geq k \geq 0$, the number of lacunar k -element subsets of $[n-1]$ is $\binom{n-k}{k}$.
- **Proof.** Write “elt” for “element”, and “subs” for “subsets”.

There is a bijection

$$\begin{aligned} \{\text{lacunar } k\text{-elt subs of } [n-1]\} &\rightarrow \{k\text{-elt subs of } \{0, 1, \dots, n-k-1\}\}, \\ \{i_1 < i_2 < \dots < i_k\} &\mapsto \{i_1 - 1 < i_2 - 2 < \dots < i_k - k\}. \end{aligned}$$

- Thus, we can start a **combinatorial proof** of our theorem as follows:

$$\sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n-k}{k} = \sum_{\substack{I \subseteq [n-1]; \\ I \text{ is lacunar}}} (-1)^{|I|}.$$

We want to prove that this is 1 or 0 or -1 .

So let us try to construct a sign-reversing involution on the set

$$\{\text{lacunar subsets of } [n-1]\}$$

except for possibly one element.

Let I be a lacunar subset of $[n-1]$.

- We try to toggle 1 in I , but we only do this if the result is lacunar.
If we succeed (i.e., if the result is lacunar), then we are done.

[Examples:

- * If $I = \{1, 3, 7\}$, then we toggle 1, and obtain the set $\{3, 7\}$. Thus, in this case, we succeed and have found the image of I under our involution.
- * If $I = \{3, 7\}$, then we toggle 1, and obtain the set $\{1, 3, 7\}$. Thus, in this case, we succeed and have found the image of I under our involution.
- * If $I = \{2, 7\}$, then we cannot toggle 1, since this would produce the non-lacunar set $\{1, 2, 7\}$. Thus, in this case, we don't succeed and move on to the next step.]
- If we have not succeeded in the previous step, then $2 \in I$ and thus $3 \notin I$.

Thus we try to toggle 4 in I , but we only do this if the result is lacunar.

If we succeed, then we are done.

[Examples:

- * If $I = \{2, 4, 9\}$, then we toggle 4, and obtain the set $\{2, 9\}$. Thus, in this case, we succeed and have found the image of I under our involution.
- * If $I = \{2, 9\}$, then we toggle 4, and obtain the set $\{2, 4, 9\}$. Thus, in this case, we succeed and have found the image of I under our involution.
- * If $I = \{2, 5, 8\}$, then we cannot toggle 4, since this would produce the non-lacunar set $\{2, 4, 5, 8\}$. Thus, in this case, we don't succeed and move on to the next step.
- * If $I = \{1, 3, 7\}$, then we do not get to this step in the first place, since the first step has already succeeded (turning I into $\{3, 7\}$).]
- If we have not succeeded in the previous step, then $5 \in I$ and thus $6 \notin I$.

Thus we try to toggle 7 in I , but we only do this if the result is lacunar.

If we succeed, then we are done.

– And so on.

This operation goes on until we have run out of elements of $[n-1]$ to toggle. The only case in which we fail to toggle anything is if


$$n \not\equiv 2 \pmod{3} \text{ and } I = \{2, 5, 8, \dots\} \cap [n-1].$$

Thus we have found a sign-reversing involution on the set $\{\text{lacunar subsets of } [n-1]\}$ with the exception of a single lacunar subset if $n \not\equiv 2 \pmod{3}$ (and with no exceptions if $n \equiv 2 \pmod{3}$). The conclusion easily follows.

- This proof is in [BenQui08] (where it is worded using domino tilings instead of lacunar subsets).

2.4. Independent sets of a graph

- Let us generalize this.
- **Definition.** Let $\Gamma = (V, E)$ be an (undirected) graph. An *independent set* of Γ means a subset I of V such that no two vertices in I are adjacent (i.e., no edge of Γ connects two vertices in I).
- **Example.** For the following graph:



the independent sets are

$$\{x, y\}, \{y, z\}, \{z, x\}, \{u, x\}, \{v, y\}, \{w, z\}, \{x, y, z\}$$

as well as all 1-element sets and the empty set.

- For any $m \geq 0$, let the *m-path* be the graph

$$1 - 2 - 3 - \dots - m$$

(that is, the graph with vertices $1, 2, \dots, m$ and edges $\{i, i+1\}$ for each $0 < i < m$).

Then, the lacunar subsets of $[m]$ are the independent sets of the *m-path*.

- Now we can generalize our previous theorem as follows:

Question: For what graphs Γ do we have

$$\sum_{\substack{I \text{ is an independent} \\ \text{set of } \Gamma}} (-1)^{|I|} \in \{1, 0, -1\} ?$$

- Certainly not for all graphs Γ (e.g., the 4-cycle is a counterexample).
- But we know it's true for path graphs. For what other graphs?
- We can try to construct a sign-reversing involution again, and see where we fail.
- What order do we try to toggle the vertices in?
- Well, we can always pick some order at random.
- Unfortunately, toggling a vertex might be blocked by several vertices.
- Trying to solve the resulting conflicts often fails (e.g., for a 4-cycle, even though the sum is -1 for a 4-cycle).
- Our above proof can be adapted when Γ is a forest ([EhrHet06, Corollary 6.1])).
- However, a much more general result holds:
- **Theorem (conjectured by Kalai and Meshulam, 1990s, proved by Chudnovsky, Scott, Seymour, Spirkl, 2018 ([CSSS18])):** Let Γ be a simple loopless undirected graph that has no induced cycle of length divisible by 3. Then,


$$\sum_{\substack{I \text{ is an independent} \\ \text{set of } \Gamma}} (-1)^{|I|} \in \{1, 0, -1\}.$$

- **Question:** Is there any proof under 10 pages length?

2.5. Dominating sets of a graph

- **Definition.** Let $\Gamma = (V, E)$ be an (undirected) graph. A *dominating set* of Γ means a subset I of V such that each vertex of Γ belongs to I or has a neighbor in I .

- **Example.** For the following graph:



the **non-dominating** sets are

$$\{u, y, z\}, \{v, z, x\}, \{w, x, y\}$$

as well as all their subsets.

- **Theorem (e.g., Brouwer 2009 ([Brouwe09], [BrCsSc09])):** The number of dominating sets of a graph Γ is always odd.
- **Theorem (Heinrich, Tittmann, 2017 ([HeiTit17], [Grinbe17, Theorem 3.2.2])):** The number of dominating sets of a graph $\Gamma = (V, E)$ is

$$2^{|V|} - 1 + \underbrace{\sum_{\substack{\text{pairs } (A,B) \text{ of disjoint} \\ \text{nonempty subsets of } V; \\ \{a,b\} \notin E \text{ for all } a \in A \text{ and } b \in B; \\ |A| \equiv |B| \pmod{2}}} (-1)^{|A|}.$$

This is even for symmetry reasons
(for any (A,B) , there is a (B,A))

- What about the alternating sum

$$\sum_{\substack{I \text{ is a dominating} \\ \text{set of } \Gamma}} (-1)^{|I|} ?$$

Is it ± 1 ?

- No; for example:

Theorem (Alikhani, 2012 ([Alikha12, Lemma 1])): If Γ is an n -cycle (for $n > 0$), then this alternating sum is

$$\begin{cases} 3, & \text{if } n \equiv 0 \pmod{4}; \\ -1, & \text{otherwise.} \end{cases}$$

Exercise: Prove this! (Is there a nice proof without too much casework?)

- **Theorem (Ehrenborg, Hetyei, 2005 ([EhrHet06, §7])):** The alternating sum is ± 1 whenever Γ is a forest.

3. Simplicial complexes

3.1. Basic definitions

- The sums we have been discussing so far didn't range over some random collections of sets. Most of them had a commonality: If a set I appeared in the sum, then so did any subset of I .
- Such collections of sets are called **simplicial complexes**.
- Formally:

Definition. A *simplicial complex* means a pair (S, Δ) , where S is a finite set and Δ is a collection (= set) of subsets of S such that

$$\text{any } I \in \Delta \text{ and } J \subseteq I \text{ satisfy } J \in \Delta.$$

- We often just write Δ for a simplicial complex (S, Δ) .
- A *face* of a simplicial complex Δ means a set $I \in \Delta$.
- Note that $\{\}$ and $\{\emptyset\}$ are two different simplicial complexes on any set S .
- **Examples** of simplicial complexes:
 - $\{\text{all subsets of } S\}$ for a given finite set S .
 - $\{\text{all lacunar subsets of } [m]\}$ for a given $m \in \mathbb{N}$.
 - $\{\text{all independent sets of } \Gamma\}$ for a given graph Γ .
 - **not** $\{\text{all dominating sets of } \Gamma\}$ for a given graph Γ .
 - $\{\text{all non-dominating sets of } \Gamma\}$ and $\{\text{all complements of dominating sets of } \Gamma\}$ for a given graph Γ .


(Here the ground set is the set of vertices of Γ .)

3.2. Geometric realizations


- Each simplicial complex (S, Δ) has a *geometric realization* $|\Delta|$, which is a topological space glued out of (geometric) simplices. The easiest way to define it is by assuming (WLOG) that $S = [n]$ for some $n \in \mathbb{N}$, and setting

$$|\Delta| = \{(t_1, t_2, \dots, t_n) \in \mathbb{R}_{\geq 0}^n \mid t_1 + t_2 + \dots + t_n = 1 \\ \text{and } \{i \mid t_i > 0\} \in \Delta\}.$$

- Normally we don't draw the literal $|\Delta|$ (since \mathbb{R}^n has too high dimension) but just something homeomorphic to it (usually in a smaller space).
- Some examples:
 - The complex $\{\text{all independent sets of } \Gamma\}$ of the graph Γ on the left is the simplicial complex drawn on the right:



- The complex $\{\text{all non-dominating sets of } \Gamma\}$ of the graph Γ on the left is the simplicial complex drawn on the right:



3.3. Homotopy and homology

- A lot of features come for free with the geometric realization:

The *homotopy type*, the *homology* and the *reduced Euler characteristic* of a simplicial complex Δ are defined to be the homotopy type, the homology and the reduced Euler characteristic of its geometric realization.

- Explicitly, the Euler characteristic of a complex Δ is simply

$$\sum_{I \in \Delta} (-1)^{|I|-1}.$$

(The “ -1 ” in the exponent just negates the whole sum.)

- Thus, the alternating sums we have been computing are actually Euler characteristics in disguise.
- Homology is a stronger invariant than Euler characteristic, and homotopy type is an even stronger invariant than homology:

$$\begin{aligned} (\text{homotopy type}) &\rightarrow (\text{homology over } \mathbb{Z}) \rightarrow (\text{homology over } \mathbb{Q}) \\ &\rightarrow (\text{Euler characteristic}). \end{aligned}$$

Our results above are all about Euler characteristics; can we lift them to those stronger invariants?

- Note that homology can be easily redefined combinatorially in terms of Δ . (Homotopy type, too, but less easily; see [Kozlov20, Proposition 9.28].)

3.4. Examples of homotopy types

- Our first theorem said that the reduced Euler characteristic of the simplicial complex

$$\{\text{all subsets of } E\}$$

is 0 for any nonempty finite set E . This lifts all the way up to homotopy level:

Proposition. This simplicial complex is contractible (i.e., homotopy-equivalent to a point).

Geometrically, this is clear: Its geometric realization is a simplex, hence homeomorphic to an $(n - 1)$ -ball, where $n = |E|$.

- Our second theorem was

$$\sum_{k=0}^m (-1)^k \binom{n}{k} = (-1)^m \binom{n-1}{m}.$$

This corresponds to the simplicial complex

$$\{\text{all subsets of } [n] \text{ having size } \leq m\}.$$

This is called the $(m - 1)$ -*skeleton* of the $(n - 1)$ -ball. By classical algebraic topology, it is homotopy-equivalent to a bouquet of $\binom{n-1}{m}$ many $(m - 1)$ -spheres, which again explains the Euler characteristic.

- Now, recall the independent sets of graphs.

Theorem (Kalai, Meshulam, Engström, Chudnovsky, Scott, Seymour, Spirk, Zhang, Wu, Kim, 2021 ([ZhaWu20], [Kim21])): Let Γ be a simple loopless undirected graph that has no induced cycle of length divisible by 3. Then, the simplicial complex

$$\{\text{independent sets of } \Gamma\}$$

is either contractible or homotopy-equivalent to a sphere (whence its reduced Euler characteristic is in $\{1, 0, -1\}$).

- As we recall, the dominating sets of a graph do not form a simplicial complex, but their complements do, and so do the non-dominating sets. As far as the alternating sum $\sum_I (-1)^{|I|}$ is concerned, these are just as good (switching between dominating and non-dominating sets or between the sets and their complements changes the sum by a factor of ± 1).

Theorem (Ehrenborg, Hetyei, 2005 ([EhrHet06, §7])): Let Γ be a forest. Then, both simplicial complexes


$$\begin{aligned} &\{\text{non-dominating sets of } \Gamma\} \quad \text{and} \\ &\{\text{complements of dominating sets of } \Gamma\} \end{aligned}$$

are either contractible or homotopy-equivalent to a sphere.

- **Question:** What can be said about the case when Γ is an n -cycle?


3.5. Discrete Morse theory

- Thus we have two approaches to proving formulas for alternating sums:



- Could these two approaches be combined? I.e., is there a technique that gets us both homotopy information and a sign-reversing involution in one

(possibly harder) swoop?



- *Discrete Morse theory* is an answer. We will use some of its very basics.
- **Definition.** For two sets I and J , we write $I \prec J$ if $J = I \cup \{\text{a single element}\}$ (that is, if $I \subseteq J$ and $|J \setminus I| = 1$). Equivalently, we write $J \succ I$ for this.
- **Definition.** Let (S, Δ) be a simplicial complex. A *partial matching* on Δ shall mean an involution $\mu : \Delta \rightarrow \Delta$ such that

$$\mu(I) = I \text{ or } \mu(I) \prec I \text{ or } \mu(I) \succ I \quad \text{for each } I \in \Delta.$$

In other words, $\mu(I)$ is either I itself or is obtained from I by removing or inserting a single element.

- **Definition.** If μ is a partial matching on Δ , then the sets $I \in \Delta$ satisfying $\mu(I) = I$ will be called *unmatched* (by μ).
- Thus, if μ is a partial matching on Δ , then

$$\sum_{I \in \Delta} (-1)^{|I|} = \sum_{\substack{I \in \Delta \\ \text{is unmatched}}} (-1)^{|I|}$$

(by cancellation).

- Thus, partial matchings are just our partial sign-reversing involutions rewritten (instead of taking some sets out of our complex, we are now leaving them fixed).
- What about the homotopy information? We cannot in general “cancel” matched faces from a simplicial complex and hope that the homotopy information is preserved.
- However, we can restrict our matchings in a way that will make them homotopy-friendly! This is one of the main contributions of Forman that became discrete Morse theory ([Forman02, §3, §6], [Kozlov20]):

- **Definition.** Let (S, Δ) be a simplicial complex. A partial matching μ on Δ is said to be *acyclic* (or a *Morse matching*) if there exists no “cycle” of the form

$$I_1 \succ \mu(I_1) \prec I_2 \succ \mu(I_2) \prec I_3 \succ \cdots \prec I_n \succ \mu(I_n) \prec I_1$$

with $n \geq 2$ and with I_1, I_2, \dots, I_n distinct.

- **Intuition:** The easiest way to ensure this is by making sure that when μ adds an element to a face I , then it does so in an “optimal” way (i.e., among all ways to add an element to I and still obtain a face of Δ , it picks the “best” one in some sense). This way, in the above “cycle”, the faces $I_1, I_2, \dots, I_n, I_1$ become “better and better”, so the cycle cannot exist. There is freedom in defining what “optimal”/“best” is (it means specifying some partial order on the faces of any given size).

This is why Forman calls acyclic matchings “*gradient vector fields*” in [Forman02].

- **Empiric fact(?):** Sign-reversing involutions in combinatorics tend to be acyclic partial matchings.
- **Question:** Really? Check some of the more complicated ones!
- **Theorem (Forman, I believe).** Let (S, Δ) be a simplicial complex, and μ an acyclic partial matching on Δ . For each $k \in \mathbb{N}$, let c_k be the number of unmatched size- k faces of Δ .

Then, there is a CW-complex homotopy-equivalent to Δ that has exactly c_k faces of dimension $k - 1$ for each $k \in \mathbb{N}$.

- **Corollary. (a)** If a simplicial complex (S, Δ) has an acyclic partial matching that leaves no face unmatched, then it is contractible.
- **(b)** If a simplicial complex (S, Δ) has an acyclic partial matching that leaves exactly one face unmatched, then it is homotopy-equivalent to a sphere.
- As a consequence, having a good Morse matching gets us good (if not 100% complete) information both about the homotopy type and about the combinatorics of a simplicial complex.
- For example, all the sign-reversing involutions we used in our proofs above are Morse matchings.


4. Elser’s “pandemic” complex

- A remarkable alternating sum identity appeared in a 1984 paper by Elser on mathematical physics (percolation theory) [Elser84]. I shall restate it in a slightly simpler language.

- Fix a (finite undirected multi)graph Γ with vertex set V and edge set E .
Fix a vertex $v \in V$.
- If $F \subseteq E$, then an F -path shall mean a path of Γ such that all edges of the path belong to F .
- If $e \in E$ is any edge and $F \subseteq E$ is any subset, then we say that F *infects* e if there exists an F -path from v to some endpoint of e .

(My go-to mental model: A virus starts out in v and spreads along any F -edge it can get to. Then, F infects e if the virus will eventually reach an endpoint of e . Note that F always infects any edge through v .)

- A subset $F \subseteq E$ is said to be *pandemic* if it infects each edge $e \in E$.
- **Example:** Let Γ be



Then:

- The set $\{1,2\} \subseteq E$ infects edges 1,2,3,6,8 (but no others), since the virus gets to the vertices v, p, q .
 - The set $\{1,2,5\}$ infects the same edges.
 - The set $\{1,2,3\}$ infects every edge other than 5.
 - The set $\{1,2,3,4\}$ infects each edge, and thus is pandemic (even though the virus never gets to vertex w).
- **Theorem (Elser, 1984 ([Elser84, Lemma 1], [Grinbe20, Theorem 1.2])):**
Assume that $E \neq \emptyset$. Then,

$$\sum_{\substack{F \subseteq E \text{ is} \\ \text{pandemic}}} (-1)^{|F|} = 0.$$


- **Remark:** A version of pandemicity in which F has to infect all vertices (rather than all edges) would fail to produce such a theorem.

4.1. More generally

- If F is a subset of E , then we define a subset $\text{Shade } F$ of E by

$$\text{Shade } F = \{e \in E \mid F \text{ infects } e\}.$$

- **Example:** Let Γ be



Then, $\text{Shade } \{1, 2\} = \{1, 2, 3, 6, 8\}$ and $\text{Shade } \{1\} = \{1, 2, 6\}$ and $\text{Shade } \{8\} = \{1, 6\}$.

- **Theorem ([Grinbe20, Theorem 2.5], generalizing Elser's theorem):** Let G be any subset of E . Assume that $E \neq \emptyset$. Then,

$$\sum_{\substack{F \subseteq E; \\ G \subseteq \text{Shade } F}} (-1)^{|F|} = 0.$$

- **Theorem ([Grinbe20, Theorem 2.6], equivalent restatement of previous theorem):** Let G be any subset of E . Then,

$$\sum_{\substack{F \subseteq E; \\ G \not\subseteq \text{Shade } F}} (-1)^{|F|} = 0.$$

- This restatement looks useful since it gets rid of the $E \neq \emptyset$ assumption. That's a good sign!

4.2. Proof idea

- Let's prove this latter restatement. Here is it again:

Theorem ([Grinbe20, Theorem 2.6], equivalent restatement of previous theorem): Let G be any subset of E . Then,

$$\sum_{\substack{F \subseteq E; \\ G \not\subseteq \text{Shade } F}} (-1)^{|F|} = 0.$$

- **Proof.** Let

$$\mathcal{A} = \{F \subseteq E \mid G \not\subseteq \text{Shade } F\}.$$

Equip the set E with a total order. If $F \in \mathcal{A}$, then let $\varepsilon(F)$ be the **smallest** edge $e \in G \setminus \text{Shade } F$.

Define a sign-reversing involution

$$\begin{aligned} \mathcal{A} &\rightarrow \mathcal{A}, \\ F &\mapsto F \triangle \{\varepsilon(F)\}. \end{aligned}$$

Check that this works! (The key observation: $\text{Shade } F$ does not change when we toggle $\varepsilon(F)$ in F .)

4.3. Variants

- We cannot replace “infects all edges” by “infects all vertices” as long as we work with sets of edges.
- However, we can work with sets of vertices instead (*mutatis mutandis*).
- In detail:
- If $F \subseteq V$, then an F -vertex-path shall mean a path of Γ such that all vertices of the path except (possibly) for its two endpoints belong to F . (Thus, if a path has only one edge or none, then it automatically is an F -vertex-path.)
- If $w \in V \setminus \{v\}$ is any vertex and $F \subseteq V \setminus \{v\}$ is any subset, then we say that F vertex-infects w if there exists an F -vertex-path from v to w . (This is always true when w is v or a neighbor of v .)
- A subset $F \subseteq V \setminus \{v\}$ is said to be *vertex-pandemic* if it vertex-infects each vertex $w \in V \setminus \{v\}$.
- **Theorem ([Grinbe20, Theorem 3.2]).** Assume that $V \setminus \{v\} \neq \emptyset$. Then,

$$\sum_{\substack{F \subseteq V \setminus \{v\} \text{ is} \\ \text{vertex-pandemic}}} (-1)^{|F|} = 0.$$

- Generalizations similar to the one above also hold.

4.4. A hammer in search of nails

- The proofs of the original Elser’s theorem and of its vertex variant are suspiciously similar.

- Even worse, they use barely any graph theory. All we needed is that E is a finite set, and that $\text{Shade} : \mathcal{P}(E) \rightarrow \mathcal{P}(E)$ (where $\mathcal{P}(E) = \{\text{all subsets of } E\}$) is a map with the property that

$$\begin{aligned} \text{Shade}(F \triangle \{u\}) &= \text{Shade } F \\ \text{for any } F \subseteq E \text{ and } u \in E \setminus \text{Shade } F. \end{aligned}$$

I call such a map Shade a *shade map*. Our above argument then shows that

$$\sum_{\substack{F \subseteq E; \\ G \not\subseteq \text{Shade } F}} (-1)^{|F|} = 0 \quad \text{for any } G \subseteq E.$$

- **Question.** Have you seen other maps satisfying this property in the wild?
- **Answer 1 ([Grinbe20, Example 4.10]).** Let A be an affine space over \mathbb{R} . Fix a finite subset E of A . For any $F \subseteq E$, we define

$$\text{Shade } F = \{e \in E \mid e \text{ is \textbf{not} a nontrivial convex combination of } F\}.$$
 (A convex combination is said to be *nontrivial* if all coefficients are < 1 .) Then, this map $\text{Shade} : \mathcal{P}(E) \rightarrow \mathcal{P}(E)$ is a shade map.
- **Answer 2 ([Grinbe20, Theorem 4.18]).** Each antimatroid (a type of greedoids, a variation on the concept of matroids) produces a shade map.
- Other answers? Can you get shade maps from matroids? spanning trees? lattices?
- **Remark ([Grinbe20, Theorem 4.33]).** Shade maps on E are in bijection with the partitions of the Boolean lattice $\mathcal{B}(E)$ into intervals.

4.5. The topological viewpoint

- Now let us return to the case of a graph $\Gamma = (V, E)$. Fix a subset G of E , and let

$$\begin{aligned} \mathcal{A} &= \{F \subseteq E \mid G \not\subseteq \text{Shade } F\} \\ &= \{F \subseteq E \mid \text{not every edge in } G \text{ is infected by } F\} \end{aligned}$$

as in the proof above.

- This \mathcal{A} is clearly a simplicial complex on ground set E .
- **Theorem (G., 2020 ([Grinbe20, Theorem 5.5])).** This simplicial complex has a Morse matching (i.e., an acyclic partial matching) with no unmatched faces. Thus, it is contractible.
- **Proof idea.** Argue that the sign-reversing involution above is a Morse matching.

4.6. The Alexander dual

- The complex

$$\mathcal{A} = \{F \subseteq E \mid G \not\subseteq \text{Shade } F\}$$

is not the only simplicial complex we can obtain from our setup. There is also

$$\mathcal{A}^* = \{F \subseteq E \mid G \subseteq \text{Shade}(E \setminus F)\}.$$

- More generally, if (S, Δ) is any simplicial complex, then we can define a new simplicial complex (S, Δ^*) , where

$$\begin{aligned} \Delta^* &:= \{I \subseteq S \mid S \setminus I \notin \Delta\} \\ &= \{\text{the complements of the non-faces of } \Delta\}. \end{aligned}$$

This (S, Δ^*) is called the *Alexander dual* of (S, Δ) .

- The homologies of (S, Δ^*) and (S, Δ) are isomorphic (folklore – see, e.g., [BjoTan09]); thus the Euler characteristics agree up to sign.
But the homotopy types are not in general equivalent! Nor is the existence of a Morse matching with good properties.
- Thus, for any homotopy type question we can answer, we can state an analogous one for its dual.
- **Question.** What is the homotopy type of the \mathcal{A}^* above?

4.7. Multi-shades?

- I can't help spreading yet another open question that essentially comes from Dorpalen-Barry et al. [DHLetc19, Conjecture 9.1].
- Return to the setup of a graph $\Gamma = (V, E)$, but don't fix the vertex v this time.
- Rename $\text{Shade } F$ as $\text{Shade}_v F$ to stress its dependence on v .
- For any subset $U \subseteq V$, define the simplicial complex

$$\mathcal{A}_U := \{F \subseteq E \mid G \not\subseteq \text{Shade}_v F \text{ for some } v \in U\}.$$

- **Question:** What can we say about the homotopy and discrete Morse theory of \mathcal{A}_U ? What about its Alexander dual?
- An optimistic yet reasonable expectation would be: a Morse matching whose unmatched faces all have the same size. (Thus, \mathcal{A}_U should be homotopy-equivalent to a bouquet of spheres.)
- **Update:** I am skeptical about the proof given in arXiv:2203.12525v2 (but I might be misunderstanding it).

5. Bonus: Path-free and path-missing complexes

- This is **joint work with Lukas Katthän and Joel Brewster Lewis** [GrKaLe21].
- Fix a **directed** graph $G = (V, E)$ and two vertices s and t . We define the two simplicial complexes

$$\mathcal{PF}(G) = \{F \subseteq E \mid \text{there is no } F\text{-path from } s \text{ to } t\}$$

(the “*path-free*” complex of G)


and

$$\mathcal{PM}(G) = \{F \subseteq E \mid \text{there is an } (E \setminus F)\text{-path from } s \text{ to } t\}$$

(the “*path-missing*” complex of G).

(These are Alexander duals of each other.)

- **Example:** Let G be the following directed graph:



Then:

- The faces of the simplicial complex $\mathcal{PF}(G)$ are the sets

$$\{b, c, e, f, g\}, \{a, c, e, f, g\}, \{b, c, d, g\}, \{a, c, d, f, g\}, \{a, b, e, f\}, \{a, b, d, f, g\}$$

as well as all their subsets.

- The faces of the simplicial complex $\mathcal{PM}(G)$ are the sets

$$\{d, e, f, g\}, \{c, d, f\}, \{a, b, c, f, g\}, \{a, e, g\}$$

as well as all their subsets.

- **Theorem (G., Katthän, Lewis, 2021 ([GrKaLe21])).** Assume that $s \neq t$ and $E \neq \emptyset$ (the other cases are trivial). Then, both complexes $\mathcal{PF}(G)$ and $\mathcal{PM}(G)$ are contractible or homotopy-equivalent to spheres. The dimensions of the spheres can be determined explicitly. The complexes are contractible if and only if G has a useless edge (i.e., an edge that appears in no path from s to t) or a (directed) cycle.

- **Theorem (G., Katthän, Lewis, 2024 ([GrKaLe21]))**. Both complexes $\mathcal{PF}(G)$ and $\mathcal{PM}(G)$ have Morse matchings with at most one unmatched face.
- The proofs use (fairly intricate) deletion/contraction arguments.
- **Question**. Is there a good non-recursive description of these Morse matchings?

Acknowledgments

Thanks to

- **Lukas Katthän**, who showed me how simple discrete Morse theory is;
- **Anders Björner, Galen Dorpalen-Barry, Dmitry Feichtner-Kozlov, Patricia Hersh, Joel Brewster Lewis, Vic Reiner, Tom Roby and Richard Stanley** for insightful conversations;
- **an anonymous referee** for one of the most useful reports I have ever gotten (within just 2 weeks), greatly simplifying one of my proofs;
- the **Mathematisches Forschungsinstitut Oberwolfach** and specifically the programme “Oberwolfach Leibniz Fellows” for its hospitality (in 2020, of all times);
- **you** for your patience!

References

- [Alikha12] Saeid Alikhani, *The Domination Polynomial of a Graph at -1* , *Graphs and Combinatorics* **29** (2013), pp. 1175–1181, doi:10.1007/s00373-012-1211-x.
- [BenQui08] Arthur T. Benjamin and Jennifer J. Quinn, *An Alternate Approach to Alternating Sums: A Method to DIE for*, *The College Mathematics Journal*, Volume 39, Number 3, May 2008, pp. 191-202(12).
- [BjoTan09] Anders Björner, Martin Tancer, *Combinatorial Alexander Duality – A Short and Elementary Proof*, *Discrete Comput Geom* (2009) 42, pp. 586–593.
- [Brouwe09] A. E. Brouwer, *The number of dominating sets of a finite graph is odd*, 2 June 2009.
<https://www.win.tue.nl/~aeb/preprints/domin2.pdf>

-
- [BrCsSc09] A. E. Brouwer, P. Csorba & A. Schrijver, *The number of dominating sets of a finite graph is odd*, 2 June 2009.
<https://www.win.tue.nl/~aeb/preprints/domin4a.pdf>
- [CSSS18] Maria Chudnovsky, Alex Scott, Paul Seymour, Sophie Spirkl, *Proof of the Kalai-Meshulam conjecture*, Israel Journal of Mathematics volume **238** (2020), pp. 639–661.
 See <https://uwspace.uwaterloo.ca/handle/10012/18516> for a preprint.
- [DHLetc19] Galen Dorpalen-Barry, Cyrus Hettle, David C. Livingston, Jeremy L. Martin, George Nasr, Julianne Vega, Hays Whitlatch, *A positivity phenomenon in Elser’s Gaussian-cluster percolation model*, arXiv:1905.11330v5, Journal of Combinatorial Theory, Series A, **179**:105364, April 2021, doi:10.1016/j.jcta.2020.105364.
- [EhrHet06] Richard Ehrenborg, Gábor Hetyei, *The topology of the independence complex*, European Journal of Combinatorics **27** (2006), pp. 906–923.
- [Elser84] Veit Elser, *Gaussian-cluster models of percolation and self-avoiding walks*, J. Phys. A: Math. Gen. **17** (1984), pp. 1515–1523.
- [Forman02] Robin Forman, *A User’s Guide to discrete Morse theory*, Séminaire Lotharingien de Combinatoire **48** (2002), Article B48c.
- [Grinbe17] Darij Grinberg, *Notes on graph theory*, 10 January 2019.
<http://www.cip.ifi.lmu.de/~grinberg/t/17s/>
- [Grinbe20] Darij Grinberg, *The Elser nuclei sum revisited*, detailed version. Also available as an ancillary file at arXiv:2009.11527v8.
- [GrKaLe21] Darij Grinberg, Lukas Kathän, Joel Brewster Lewis, *The path-missing and path-free complexes of a directed graph*, arXiv:2102.0784v3.
- [HeiTit17] Irene Heinrich, Peter Tittmann, *Counting Dominating Sets of Graphs*, arXiv:1701.03453v1.
- [Kim21] Jinha Kim, *The homotopy type of the independence complex of graphs with no induced cycle of length divisible by 3*, arXiv:2101.07131v3.
- [Kozlov20] Dmitry N. Kozlov, *Organized Collapse: An Introduction to Discrete Morse Theory*, Graduate Studies in Mathematics **207**, AMS 2020.
 See <https://dfklab.com/books/> for a preprint.
- [Sagan20] Bruce Sagan, *Combinatorics: The Art of Counting*, Graduate Studies in Mathematics **210**, AMS 2020.
 See <https://users.math.msu.edu/users/bsagan/Books/Aoc/final.pdf> for a preprint.
-

- [Stanle11] Richard P. Stanley, *Enumerative Combinatorics, volume 1*, 2nd edition, version of 15 July 2011.
<http://math.mit.edu/~rstan/ec/> .
See <http://math.mit.edu/~rstan/ec/> for errata.
- [Striker15] Jessica Striker, *The toggle group, homomesy, and the Razumov-Stroganov correspondence*, arXiv:1503.08898v2, Electron. J. Combin. **22** (2015) no. 2.
- [ZhaWu20] Hehui Wu, Wentao Zhang, *The Betti Number of the Independence Complex of Ternary Graphs*, arXiv:2011.10939v3.
-