

Math 530: Graph Theory, Spring 2023:  
Homework 9  
due 2023-06-11 at 11:59 PM  
**Please solve 3 of the 9 problems!**

---

Darij Grinberg

July 23, 2023

---

1 EXERCISE 1

1.1 PROBLEM

Let  $D = (V, A, \psi)$  be a strongly connected multidigraph.

A *wealth distribution* on  $D$  shall mean a family  $(k_v)_{v \in V}$  of integers (one for each vertex  $v \in V$ ). If  $k = (k_v)_{v \in V}$  is a wealth distribution, then we refer to each value  $k_v$  as the *wealth* of the vertex  $v$ , and we define the *total wealth* of  $k$  to be the sum  $\sum_{v \in V} k_v$ . We say that a vertex  $v$  is *in debt* in a given wealth distribution  $k = (k_v)_{v \in V}$  if its wealth  $k_v$  is negative.


For any vertices  $v$  and  $w$ , we let  $a_{v,w}$  denote the number of arcs that have source  $v$  and  $w$ .

A *donation* is an operation that transforms a wealth distribution as follows: We choose a vertex  $v$ , and we decrease its wealth by its outdegree  $\deg^+ v$ , and then increase the wealth of each vertex  $w \in V$  (including  $v$  itself) by  $a_{v,w}$ . (You can think of  $v$  as donating a unit of wealth for each arc that has source  $v$ . This unit flows to the target to this arc. Note that a donation does not change the total wealth.)

Let  $k$  be a wealth distribution on  $D$  whose total wealth is larger than  $|A| - |V|$ . Prove that by an appropriately chosen finite sequence of donations, we can ensure that no vertex is in debt.

## 1.2 REMARK

For instance, consider the digraph



with wealth distribution  $(k_1, k_2, k_3, k_4, k_5, k_6) = (-1, -1, 1, 2, 0, 1)$ . The vertices 1 and 2 are in debt here, but it is possible to get all vertices out of debt by having the vertices 4, 5, 6, 1 donate in some order (the order clearly does not matter for the result<sup>1</sup>).

Note that vertices are allowed to donate multiple times (although in the above example, this was unnecessary).

## 1.3 HINT

Show first that if the total wealth is larger than  $|A| - |V|$ , then at least one vertex  $v$  has wealth  $\geq \deg^+ v$ .

## 1.4 SOLUTION

...

---

## 2 EXERCISE 2

## 2.1 PROBLEM

We continue with the setting and terminology of Exercise 1.

A *clawback* is an operation that transforms a wealth distribution as follows: We choose a vertex  $v$ , and we increase its wealth by its outdegree  $\deg^+ v$ , and then decrease the wealth of each vertex  $w \in V$  (including  $v$  itself) by  $a_{v,w}$ . (Thus, a clawback is the inverse of a donation.)


Let  $k$  be a wealth distribution on  $D$  whose total wealth is larger than  $|A| - |V|$ . Prove that by an appropriately chosen finite sequence of clawbacks, we can ensure that no vertex is in debt.

---

<sup>1</sup>Depending on the order, some vertices will go into debt in the process, but this is okay as long as they ultimately end up debt-free.

## 2.2 REMARK

Note that we are still assuming  $D$  to be strongly connected. Otherwise, the truth of the claim is not guaranteed. For instance, for the digraph



with wealth distribution  $(k_1, k_2, k_3, k_4) = (0, 0, -1, 2)$ , no sequence of donations and clawbacks will result in every vertex being out of debt (since the wealth difference  $k_4 - k_3$  is preserved under any donation or clawback, but this difference is too large to come from a debt-free distribution with total weight 1).

## 2.3 HINT

Show that any donation is equivalent to an appropriately chosen composition of clawbacks.

## 2.4 SOLUTION

...

---

## 3 EXERCISE 3

## 3.1 PROBLEM

Let  $X$  and  $Y$  be two finite sets such that  $|X| \leq |Y|$ . Let  $f : X \rightarrow Y$  be a map that is not constant. (A map is said to be *constant* if all its values are equal.) Prove that there exists an injective map  $g : X \rightarrow Y$  such that each  $x \in X$  satisfies  $g(x) \neq f(x)$ .

## 3.2 SOLUTION

...

---

## 4 EXERCISE 4

## 4.1 PROBLEM

Let  $(G, X, Y)$  be a bipartite graph with  $X \neq \emptyset$ . Assume that  $G$  has an  $X$ -complete matching.

An edge  $e$  of  $G$  will be called *useless* if  $G$  has no  $X$ -complete matching that contains  $e$ . Prove that there exists a vertex  $x \in X$  such that no edge that contains  $x$  is useless.

## 4.2 SOLUTION

...

## 5 EXERCISE 5

## 5.1 PROBLEM

Let  $G = (V, E, \varphi)$  be a multigraph. Let  $M$  be a matching of  $G$ .

An *augmenting path* for  $M$  shall mean a path  $(v_0, e_1, v_1, e_2, v_2, \dots, e_k, v_k)$  of  $G$  such that  $k$  is odd<sup>2</sup> and such that

- the even-indexed edges  $e_2, e_4, \dots, e_{k-1}$  belong to  $M$  (note that this condition is vacuously true if  $k = 1$ );
- the odd-indexed edges  $e_1, e_3, \dots, e_k$  belong to  $E \setminus M$ ;
- neither the starting point  $v_0$  nor the ending point  $v_k$  is matched in  $M$ .

Prove that  $M$  has maximum size among all matchings of  $G$  if and only if there exists no augmenting path for  $M$ .

## 5.2 HINT

If  $M$  and  $M'$  are two matchings of  $G$ , what can you say about the symmetric difference  $(M \cup M') \setminus (M \cap M')$ ?

## 5.3 SOLUTION

...

## 6 EXERCISE 6

## 6.1 PROBLEM

Let  $(G, X, Y)$  be a bipartite graph. Prove that

$$\sum_{A \subseteq X} (-1)^{|A|} [N(A) = Y] = \sum_{B \subseteq Y} (-1)^{|B|} [N(B) = X]$$

(where we are using the Iverson bracket notation).

<sup>2</sup>Note that  $k = 1$  is allowed.

## 6.2 SOLUTION

...

## 7 EXERCISE 7

## 7.1 PROBLEM

Let  $A$  and  $B$  be two finite sets such that  $|B| \geq |A|$ . Let  $d_{i,j}$  be a real number for each  $(i, j) \in A \times B$ . Let<sup>3</sup>

$$m_1 = \min_{\sigma: A \rightarrow B \text{ injective}} \max_{i \in A} d_{i, \sigma(i)}$$

and

$$m_2 = \max_{\substack{I \subseteq A; J \subseteq B; \\ |I| + |J| = |B| + 1}} \min_{(i,j) \in I \times J} d_{i,j}.$$

Prove that  $m_1 = m_2$ .

## 7.2 SOLUTION

...

## 8 EXERCISE 8

## 8.1 PROBLEM

Let  $c$  and  $r$  be two positive integers. Let  $T$  be a tournament with more than  $r^c$  vertices. Each arc of  $T$  is colored with one of the  $c$  colors  $1, 2, \dots, c$ . Prove that  $T$  has a monochromatic path of length  $r$ .

(A path is said to be *monochromatic* if all its arcs have the same color.)

## 8.2 HINT

Induct on  $c$ , and apply Gallai-Milgram to a certain digraph in the induction step. (Note that the case  $c = 1$  recovers the Easy Redei Theorem.)

## 8.3 SOLUTION

...

<sup>3</sup>The notation “ $\min_{\text{some kind of objects}} \text{some kind of value}$ ” means the minimum of the given value over all objects of the given kind. An analogous notation is used for a maximum.

## 9 EXERCISE 9

### 9.1 PROBLEM

Let  $D$  be a balanced multidigraph. Let  $s$  and  $t$  be two vertices of  $D$ . Let  $k \in \mathbb{N}$ . Assume that  $D$  has  $k$  pairwise arc-disjoint paths from  $s$  to  $t$ . Show that  $D$  has  $k$  pairwise arc-disjoint paths from  $t$  to  $s$ .

[We say that  $k$  paths  $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$  are *pairwise arc-disjoint* if and only if no arc appears in more than one of these  $k$  paths.]

### 9.2 SOLUTION

...

---