Drexel University, Department of Mathematics

Math 332: Undergraduate Abstract Algebra II, Winter 2023: Midterm 1

Please solve **at most 3 of the 6 problems!** **No collaboration** is allowed on the midterm.

Darij Grinberg

February 26, 2023

1 Exercise 1

1.1 Problem

(a) Is the map

$$\mathbb{Z}^{2\times 2} \to \mathbb{Z}^{2\times 2},$$
$$A \mapsto A^T$$

(which sends each 2×2-matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ to its transpose $A^T = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$) a ring morphism?

(b) Is the map

$$\mathbb{Z}^{2\times 2} \to \mathbb{Z}^{2\times 2},$$
$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} d & c \\ b & a \end{pmatrix}$$

a ring morphism?

Keep in mind that claims should be proved.

1.2 Solution

...

---

2 Exercise 2

2.1 Problem

Let R be a ring. Let a and b be two units of R such that $a+b$ is a unit as well.

(a) Prove that $a^{-1}+b^{-1}$, too, is a unit, and its inverse is

$$\left(a^{-1}+b^{-1}\right)^{-1} = a \cdot (a+b)^{-1} \cdot b = b \cdot (a+b)^{-1} \cdot a.$$

(b) Show on an example that $\left(a^{-1}+b^{-1}\right)^{-1}$ can differ from $ab \cdot (a+b)^{-1}$.

2.2 Solution

...

---

3 Exercise 3

3.1 Problem

Let R be a ring. If A, B, C, D are four subsets of R, then the notation $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ shall denote the set of all 2×2-matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in R^{2 \times 2}$ with $a \in A$, $b \in B$, $c \in C$ and $d \in D$. (For instance, $\begin{pmatrix} \mathbb{N} & 2\mathbb{Z} \\ 2\mathbb{Z} & \mathbb{N} \end{pmatrix}$ is the set of all 2×2-matrices whose diagonal entries are nonnegative integers and whose off-diagonal entries are even integers.)

(a) Let I be a subset of R. Prove that I is an ideal of R if and only if $\begin{pmatrix} R & I \\ \{0\} & R \end{pmatrix}$ is a subring of $R^{2 \times 2}$.

(b) Does the same claim hold for $\begin{pmatrix} R & I \\ I & R \end{pmatrix}$ instead of $\begin{pmatrix} R & I \\ \{0\} & R \end{pmatrix}$?

(c) Does the same claim hold for $\begin{pmatrix} R & I \\ R & R \end{pmatrix}$ instead of $\begin{pmatrix} R & I \\ \{0\} & R \end{pmatrix}$?

(d) Does the same claim hold for $\begin{pmatrix} R & R \\ R & I \end{pmatrix}$ instead of $\begin{pmatrix} R & I \\ \{0\} & R \end{pmatrix}$?

For the sake of brevity, you need not give proofs for parts (b), (c) and (d).

3.2 Solution

...

---

4 Exercise 4

4.1 Problem

Let R be a commutative ring in which $8 \cdot 1_R = 0_R$. (Examples of such rings are $\mathbb{Z}/2$, $\mathbb{Z}/4$ and $\mathbb{Z}/8$, but there are also many others, such as polynomial rings over $\mathbb{Z}/8$.)

Prove that the set of all elements $a \in R$ satisfying $(1-2a)^2 = 1$ is a subring of R.

4.2 Solution

...

---

5 Exercise 5

5.1 Problem

Consider the ring $\mathbb{R}^{2\times 2}$ of all 2×2-matrices with real entries.

Define two subsets $\mathcal{P}$ and $\mathcal{M}$ of $\mathbb{R}^{2\times 2}$ by

$$\mathcal{P} := \left\{ \begin{pmatrix} a & b \\ b & a \end{pmatrix} \mid a, b \in \mathbb{R} \right\} \qquad \text{and}$$
$$\mathcal{M} := \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \mid a, b \in \mathbb{R} \right\}.$$

(a) Show that $\mathcal{P}$ and $\mathcal{M}$ are commutative subrings of $\mathbb{R}^{2\times 2}$.

(b) Prove that $\mathcal{P}$ is not an integral domain.

(c) Prove that $\mathcal{M}$ is a field.

5.2 Solution

...

---

6 Exercise 6

6.1 Problem

Let R be a ring. For any two subsets A and B of R, we define a subset $A+B$ of R by

$$A+B := \{a+b \mid a \in A \text{ and } b \in B\}.$$

Which of the following three claims are true? (Prove the true ones and give counterexamples to the false ones.)

(a) If I and J are two ideals of R, then $I+J$ is again an ideal of R.

(b) If A and B are two subrings of R, then $A+B$ is again a subring of R.

(c) If I is an ideal of R, and if S is a subring of R, then $S+I$ is a subring of R.

6.2 Solution

...