

Math 221 Winter 2023 (Darij Grinberg): homework set 5

due date: Sunday 2023-03-13 at 11:59PM on gradescope (



<https://www.gradescope.com/courses/487830>).Please solve only **4 of the 6 exercises**.

The first two exercises are about functions and their properties:

Exercise 1. Let A, B, C, D be four sets. Let $f : A \rightarrow C$ and $g : B \rightarrow D$ be two maps. Define a new map $f * g : A \times B \rightarrow C \times D$ by setting

$$(f * g)(a, b) = (f(a), g(b)) \quad \text{for every pair } (a, b) \in A \times B.$$

Prove the following:

(a) If f and g are injective, then $f * g$ is injective.**(b)** If f and g are surjective, then $f * g$ is surjective.**Exercise 2.** Let X and Y be two sets. Let $f : X \rightarrow Y$ be a map.A **left inverse** of f means a map $g : Y \rightarrow X$ that satisfies $g \circ f = \text{id}_X$ (but not necessarily $f \circ g = \text{id}_Y$).A **right inverse** of f means a map $g : Y \rightarrow X$ that satisfies $f \circ g = \text{id}_Y$ (but not necessarily $g \circ f = \text{id}_X$).**(a)** Prove that f has a right inverse if and only if f is surjective.**(b)** Assume that $X \neq \emptyset$. Prove that f has a left inverse if and only if f is injective.**(c)** Find two distinct left inverses of the map**(d)** Find two distinct right inverses of the map

Some exercises on counting follow:

Exercise 3. Let $n \in \mathbb{N}$. Compute the # of 4-tuples $(a, b, c, d) \in [n]^4$ that satisfy $a \leq b < c \leq d$. (Not a typo: the second sign is a $<$, not a $\leq$.)

(Recall that $[n]^4 = [n] \times [n] \times [n] \times [n]$, so that a 4-tuple $(a, b, c, d) \in [n]^4$ means a 4-tuple of integers $a, b, c, d \in \{1, 2, \dots, n\}$.)

Exercise 4. Let $n \in \mathbb{N}$. Compute the # of pairs (A, B) of subsets of $[n]$ that satisfy $A \cap B = \emptyset$.

(For example, if $n = 2$, then this # is 9, since there are 9 such pairs:

$$\begin{aligned} &(\emptyset, \emptyset), \quad (\emptyset, \{1\}), \quad (\emptyset, \{2\}), \quad (\emptyset, \{1, 2\}), \\ &(\{1\}, \emptyset), \quad (\{1\}, \{2\}), \quad (\{2\}, \emptyset), \quad (\{2\}, \{1\}), \\ &(\{1, 2\}, \emptyset). \end{aligned}$$

)

Exercise 5. A set S of integers will be called **pseudolacunar** if it has the property that no two elements s, t of S satisfy $|s - t| = 2$. For instance, the set $\{2, 5, 6\}$ is pseudolacunar, but the set $\{2, 5, 7\}$ is not (since $|5 - 7| = 2$).

For each $n \in \mathbb{N}$, let p_n be the # of pseudolacunar subsets of $[n]$.

Prove that

$$p_n = p_{n-1} + p_{n-3} + p_{n-4} \quad \text{for each } n \geq 4.$$

[Hint: To each pseudolacunar subset, assign one of three colors.]

Exercise 6. A set S of integers shall be called **self-starting** if its size $|S|$ is also its smallest element. (For example, $\{3, 5, 6\}$ is self-starting, while $\{2, 3, 4\}$ and $\{3\}$ are not.)

Let $n \in \mathbb{N}$.

(a) For any $k \in [n]$, find the number of self-starting subsets of $[n]$ having size k .

(b) Find the number of all self-starting subsets of $[n]$.