

# From Chio Pivotal Condensation to the Matrix-Tree theorem

Darij Grinberg, Karthik Karnik, Anya Zhang

January 10, 2019

## Abstract

We show a determinant identity which generalizes both the Chio pivotal condensation theorem and the Matrix-Tree theorem.

## 1. Introduction

The Chio pivotal condensation theorem (Theorem 2.1 below, or [Eves68, Theorem 3.6.1]) is a simple particular case of the Dodgson-Muir determinantal identity ([BerBru08, (4)]), which can be used to reduce the computation of an  $n \times n$ -determinant to that of an  $(n - 1) \times (n - 1)$ -determinant (provided that an entry of the matrix can be divided by<sup>1</sup>). On the other hand, the Matrix-Tree theorem (Theorem 2.12, or [Zeilbe85, Section 4], or [Verstr12, Theorem 1]) expresses the number of spanning trees of a graph as a determinant<sup>2</sup>. In this note, we show that these two results have a common generalization (Theorem 2.13). As we have tried to keep the note self-contained, using only the well-known fundamental properties of determinants, it also provides new proofs for both results.

### 1.1. Acknowledgments

We thank the PRIMES project at MIT, during whose 2015 iteration this paper was created, and in particular George Lusztig for sponsoring the first author's mentorship in this project.

---

<sup>1</sup>We work with matrices over arbitrary commutative rings, so this is not a moot point. Of course, if the ring is a field, then this just means that the matrix has a nonzero entry.

<sup>2</sup>And not just the number; rather, a "weighted number" from which the spanning trees can be read off if the weights are chosen generically enough.

## 2. The theorems

We shall use the (rather standard) notations defined in [Grinbe15]. In particular,  $\mathbb{N}$  means the set  $\{0, 1, 2, \dots\}$ . For any  $n \in \mathbb{N}$ , we let  $S_n$  denote the group of permutations of the set  $\{1, 2, \dots, n\}$ . The  $n \times m$ -matrix whose  $(i, j)$ -th entry is  $a_{i,j}$  for each  $(i, j) \in \{1, 2, \dots, n\} \times \{1, 2, \dots, m\}$  will be denoted by  $(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq m}$ .

Let  $\mathbb{K}$  be a commutative ring. We shall regard  $\mathbb{K}$  as fixed throughout this note (so we won't always write "Let  $\mathbb{K}$  be a commutative ring" in our propositions); the notion "matrix" will always mean "matrix with entries in  $\mathbb{K}$ ".

### 2.1. Chio Pivotal Condensation

We begin with a statement of the Chio Pivotal Condensation theorem (see, e.g., [KarZha16, Theorem 0.1] and the reference therein):

**Theorem 2.1.** Let  $n \geq 2$  be an integer. Let  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  be a matrix. Then,

$$\det \left( (a_{i,j}a_{n,n} - a_{i,n}a_{n,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) = a_{n,n}^{n-2} \cdot \det \left( (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \right).$$

**Example 2.2.** If  $n = 3$  and  $A = \begin{pmatrix} a & a' & a'' \\ b & b' & b'' \\ c & c' & c'' \end{pmatrix}$ , then Theorem 2.1 says that

$$\det \begin{pmatrix} ac'' - a''c & a'c'' - a''c' \\ bc'' - b''c & b'c'' - b''c' \end{pmatrix} = (c'')^{3-2} \cdot \det \begin{pmatrix} a & a' & a'' \\ b & b' & b'' \\ c & c' & c'' \end{pmatrix}.$$

Theorem 2.1 (originally due to Félix Chio in 1853<sup>3</sup>) is nowadays usually regarded either as a particular case of the Dodgson-Muir determinantal identity ([BerBru08, (4)]), or as a relatively easy exercise on row operations and the method of universal identities<sup>4</sup>. We, however, shall generalize it in a different direction.

<sup>3</sup>See [Heinig11, footnote 2] and [Abeles14, §2] for some historical background.

<sup>4</sup>In more detail:

- In order to derive Theorem 2.1 from [BerBru08, (4)], it suffices to set  $k = n - 1$  and recognize the right hand side of [BerBru08, (4)] as  $\det \left( (a_{i,j}a_{n,n} - a_{i,n}a_{n,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right)$ .
- A proof of Theorem 2.1 using row operations can be found in [Eves68, Theorem 3.6.1], up to a few minor issues: First of all, [Eves68, Theorem 3.6.1] proves not exactly Theorem 2.1 but the analogous identity

$$\det \left( (a_{i+1,j+1}a_{1,1} - a_{i+1,1}a_{1,j+1})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) = a_{1,1}^{n-2} \cdot \det \left( (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \right).$$

## 2.2. Generalization, step 1

Our generalization will proceed in two steps. In the first step, we shall replace some of the  $n$ 's on the left hand side by  $f(i)$ 's (see Theorem 2.9 below). We first define some notations:

**Definition 2.3.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be any map such that  $f(n) = n$ .

We say that the map  $f$  is  $n$ -potent if for every  $i \in \{1, 2, \dots, n\}$ , there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$ . (In less formal terms,  $f$  is  $n$ -potent if and only if every element of  $\{1, 2, \dots, n\}$  eventually arrives at  $n$  when being subjected to repeated application of  $f$ .)

(Note that, by definition, any  $n$ -potent map  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  must satisfy  $f(n) = n$ .)

**Example 2.4.** For this example, let  $n = 3$ . The map  $\{1, 2, 3\} \rightarrow \{1, 2, 3\}$  sending  $1, 2, 3$  to  $2, 1, 3$ , respectively, is not  $n$ -potent (because applying it repeatedly to 1 can only give 1 or 2, but never 3). The map  $\{1, 2, 3\} \rightarrow \{1, 2, 3\}$  sending  $1, 2, 3$  to  $3, 3, 2$ , respectively, is not  $n$ -potent (since it does not send  $n$  to  $n$ ). The map  $\{1, 2, 3\} \rightarrow \{1, 2, 3\}$  sending  $1, 2, 3$  to  $3, 1, 3$ , respectively, is  $n$ -potent (indeed, every element of  $\{1, 2, 3\}$  goes to 3 after at most two applications of this map).

**Remark 2.5.** Given a positive integer  $n$ , the  $n$ -potent maps  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  are in 1-to-1 correspondence with the trees with vertex set  $\{1, 2, \dots, n\}$ . Namely, an  $n$ -potent map  $f$  corresponds to the tree whose edges are  $\{i, f(i)\}$  for all  $i \in \{1, 2, \dots, n-1\}$ . If we regard the tree as a rooted tree with root  $n$ , and if we direct every edge towards the root, then the edges are  $(i, f(i))$  for all  $i \in \{1, 2, \dots, n-1\}$ .

**Remark 2.6.** Let  $n \geq 2$  be an integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be any  $n$ -potent map. Then:

- (a) There exists some  $g \in \{1, 2, \dots, n-1\}$  such that  $f(g) = n$ .
- (b) We have  $|f^{-1}(n)| \geq 2$ .

The (very simple) proof of Remark 2.6 can be found in the Appendix (Section 4).

---

Second, [Eves68, Theorem 3.6.1] assumes  $a_{1,1}$  to be invertible (and all  $a_{i,j}$  to belong to a field); however, assumptions like this can easily be disposed of using the method of universal identities (see [Conrad09]).

A more explicit and self-contained proof of Theorem 2.1 can be found in [KarZha16]. References to other proofs appear in [Abeles14, §2].

---

**Definition 2.7.** Let  $n \geq 2$  be an integer. Let  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  be an  $n \times n$ -matrix. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be any  $n$ -potent map.

(a) We define an element  $\text{weight}_f A$  of  $\mathbb{K}$  by

$$\text{weight}_f A = \prod_{i=1}^{n-1} a_{i,f(i)}.$$

(b) We define an element  $\text{abut}_f A$  of  $\mathbb{K}$  by

$$\text{abut}_f A = a_{n,n}^{|f^{-1}(n)|-2} \prod_{\substack{i \in \{1, 2, \dots, n-1\}; \\ f(i) \neq n}} a_{f(i),n}.$$

(This is well-defined, since Remark 2.6 (b) shows that  $|f^{-1}(n)| - 2 \in \mathbb{N}$ .)

**Remark 2.8.** Let  $n$ ,  $A$  and  $f$  be as in Definition 2.7. Here are two slightly more intuitive ways to think of  $\text{abut}_f A$ :

(a) If  $a_{n,n} \in \mathbb{K}$  is invertible, then  $\text{abut}_f A$  is simply  $\frac{1}{a_{n,n}} \prod_{i \in \{1, 2, \dots, n-1\}} a_{f(i),n}$ .

(b) Remark 2.6 (a) shows that there exists some  $g \in \{1, 2, \dots, n-1\}$  such that  $f(g) = n$ . Fix such a  $g$ . Then,

$$\text{abut}_f A = \prod_{\substack{i \in \{1, 2, \dots, n-1\}; \\ i \neq g}} a_{f(i),n}.$$

The (nearly trivial) proof of Remark 2.8 is again found in the Appendix. Now, we can state our first generalization of Theorem 2.1:

**Theorem 2.9.** Let  $n$  be a positive integer. Let  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  be an  $n \times n$ -matrix. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be any map such that  $f(n) = n$ . Let  $B$  be the  $(n-1) \times (n-1)$ -matrix

$$\left( a_{i,j} a_{f(i),n} - a_{i,n} a_{f(i),j} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \in \mathbb{K}^{(n-1) \times (n-1)}.$$

(a) If the map  $f$  is not  $n$ -potent, then  $\det B = 0$ .

(b) Assume that  $n \geq 2$ . Assume that the map  $f$  is  $n$ -potent. Then,

$$\det B = (\text{abut}_f A) \cdot \det A.$$

**Example 2.10.** For this example, let  $n = 3$  and  $A = \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \end{pmatrix}$ .

If  $f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$  is the map sending 1, 2, 3 to 3, 1, 3, respectively, then the matrix  $B$  defined in Theorem 2.9 is  $\begin{pmatrix} a_{1,1}a_{3,3} - a_{1,3}a_{3,1} & a_{1,2}a_{3,3} - a_{1,3}a_{3,2} \\ a_{2,1}a_{1,3} - a_{2,3}a_{1,1} & a_{2,2}a_{1,3} - a_{2,3}a_{1,2} \end{pmatrix}$ . Since this map  $f$  is  $n$ -potent, Theorem 2.9 (b) predicts that this matrix  $B$  satisfies  $\det B = (\text{abut}_f A) \cdot \det A$ . This is indeed easily checked (indeed, we have  $\text{abut}_f A = a_{1,3}$  in this case).

On the other hand, if  $f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$  is the map sending 1, 2, 3 to 1, 1, 3, respectively, then the matrix  $B$  defined in Theorem 2.9 is  $\begin{pmatrix} a_{1,1}a_{1,3} - a_{1,3}a_{1,1} & a_{1,2}a_{1,3} - a_{1,3}a_{1,2} \\ a_{2,1}a_{1,3} - a_{2,3}a_{1,1} & a_{2,2}a_{1,3} - a_{2,3}a_{1,2} \end{pmatrix}$ . Since this map  $f$  is not  $n$ -potent, Theorem 2.9 (a) predicts that this matrix  $B$  satisfies  $\det B = 0$ . This, too, is easily checked (and arguably obvious in this case).

Applying Theorem 2.9 (b) to  $f(i) = n$  yields Theorem 2.1. (The map  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  defined by  $f(i) = n$  is clearly  $n$ -potent, and satisfies  $\text{abut}_f A = a_{n,n}^{n-2}$ .)

We defer the proof of Theorem 2.9 until later; first, let us see how it can be generalized a bit further (not substantially, anymore) and how this generalization also encompasses the matrix-tree theorem.

### 2.3. The matrix-tree theorem

**Definition 2.11.** For any two objects  $i$  and  $j$ , we define an element  $\delta_{i,j} \in \mathbb{K}$  by  $\delta_{i,j} = \begin{cases} 1, & \text{if } i = j; \\ 0, & \text{if } i \neq j. \end{cases}$

Let us first state the matrix-tree theorem.

To be honest, there is no “the matrix-tree theorem”, but rather a network of “matrix-tree theorems” (some less, some more general), each of which has a reasonable claim to this name. Here we shall prove the following one:

**Theorem 2.12.** Let  $n \geq 1$  be an integer. Let  $W : \{1, 2, \dots, n\} \times \{1, 2, \dots, n\} \rightarrow \mathbb{K}$  be any function. For every  $i \in \{1, 2, \dots, n\}$ , set

$$d^+(i) = \sum_{j=1}^n W(i, j).$$

Let  $L$  be the matrix  $(\delta_{i,j}d^+(i) - W(i, j))_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \in \mathbb{K}^{(n-1) \times (n-1)}$ . Then,

$$\det L = \sum_{\substack{f: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}; \\ f(n) = n; \\ f \text{ is } n\text{-potent}}} \prod_{i=1}^{n-1} W(i, f(i)). \tag{1}$$

Since our notation differs from that in most other sources on the matrix-tree theorem, let us explain the equivalence between our Theorem 2.12 and one of its better-known avatars: The version of the matrix-tree theorem stated in [Zeilbe85, Section 4] involves some “weights”  $a_{k,m}$ , a determinant of an  $(n-1) \times (n-1)$ -matrix, and a sum over a set  $\mathcal{T} = \mathcal{T}(n)$ . These correspond (respectively) to the values  $W(k,m)$ , the determinant  $\det L$ , and the sum over all  $n$ -potent maps  $f$  in our Theorem 2.12. In fact, the only nontrivial part of this correspondence is the bijection between the trees in  $\mathcal{T}$  and the  $n$ -potent maps  $f$  over which the sum in (1) ranges. This bijection is precisely the one introduced in Remark 2.5.<sup>5</sup>

It might seem weird to call Theorem 2.12 the “matrix-tree theorem” if the word “tree” never occurs inside it. However, as we have already noticed in Remark 2.5, the trees on the set  $\{1, 2, \dots, n\}$  are in bijection with the  $n$ -potent maps  $\{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ , and therefore the sum on the right hand side of (1) can be viewed as a sum over all these trees. Moreover, the function  $W$  can be viewed as an  $n \times n$ -matrix; when this matrix is specialized to the adjacency matrix of a directed graph, the sum on the right hand side of (1) becomes the number of directed spanning trees of this directed graph directed towards the root  $n$ .

## 2.4. Generalization, step 2

Now, as promised, we will generalize Theorem 2.9 a step further. While the result will not be significantly stronger (we will actually derive it from Theorem 2.9 quite easily), it will lead to a short proof of Theorem 2.12:

---

<sup>5</sup>A slightly different version of the matrix-tree theorem appears in [Verstr12, Theorem 1] (and various other places); it involves a function  $W$ , a number  $v \in \{1, 2, \dots, n\}$ , a matrix  $L_v$ , a set  $\mathcal{T}_v$  and a sum  $\tau(W, v)$ . Our Theorem 2.12 is equivalent to the case of [Verstr12, Theorem 1] for  $v = n$ ; but this case is easily seen to be equivalent to the general case of [Verstr12, Theorem 1] (since the elements of  $\{1, 2, \dots, n\}$  can be permuted at will). Our matrix  $L$  is the  $L_n$  of [Verstr12, Theorem 1]. Furthermore, our sum over all  $n$ -potent maps  $f$  corresponds to the sum  $\tau(W, n)$  in [Verstr12], which is a sum over all  $n$ -arborescences on  $\{1, 2, \dots, n\}$ ; the correspondence is again due to Remark 2.5.

---

**Theorem 2.13.** Let  $n \geq 2$  be an integer. Let  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  and  $B = (b_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  be  $n \times n$ -matrices. Write the  $n \times n$ -matrix  $BA$  in the form  $BA = (c_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ .

Let  $G$  be the  $(n - 1) \times (n - 1)$ -matrix

$$(a_{i,j}c_{i,n} - a_{i,n}c_{i,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \in \mathbb{K}^{(n-1) \times (n-1)}.$$

Then,

$$\det G = \left( \sum_{\substack{f: \{1,2,\dots,n\} \rightarrow \{1,2,\dots,n\}; \\ f(n)=n; \\ f \text{ is } n\text{-potent}}} (\text{weight}_f B) (\text{abut}_f A) \right) \cdot \det A.$$

To obtain Theorem 2.9 from Theorem 2.13, we have to define  $B$  by  $B = (\delta_{j,f(i)})_{1 \leq i \leq n, 1 \leq j \leq n}$ . Below we shall show how to obtain the matrix-tree theorem from Theorem 2.13.

**Example 2.14.** Let us see what Theorem 2.13 says for  $n = 3$ . There are three  $n$ -potent maps  $f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$ :

- one map  $f_{33}$  which sends both 1 and 2 to 3;
- one map  $f_{23}$  which sends 1 to 2 and 2 to 3;
- one map  $f_{31}$  which sends 2 to 1 and 1 to 3.

The definition of the  $c_{i,j}$  as the entries of  $BA$  shows that  $c_{i,j} = b_{i,1}a_{1,j} + b_{i,2}a_{2,j} + b_{i,3}a_{3,j}$  for all  $i$  and  $j$ . We have

$$G = \begin{pmatrix} a_{1,1}c_{1,3} - c_{1,1}a_{1,3} & a_{1,2}c_{1,3} - c_{1,2}a_{1,3} \\ a_{2,1}c_{2,3} - c_{2,1}a_{2,3} & a_{2,2}c_{2,3} - c_{2,2}a_{2,3} \end{pmatrix}.$$

Theorem 2.13 says that

$$\begin{aligned} \det G &= \left( (\text{weight}_{f_{33}} B) (\text{abut}_{f_{33}} A) + (\text{weight}_{f_{23}} B) (\text{abut}_{f_{23}} A) \right. \\ &\quad \left. + (\text{weight}_{f_{31}} B) (\text{abut}_{f_{31}} A) \right) \cdot \det A \\ &= (b_{1,3}b_{2,3}a_{3,3} + b_{1,2}b_{2,3}a_{2,3} + b_{1,3}b_{2,1}a_{1,3}) \cdot \det A. \end{aligned}$$

### 3. The proofs

#### 3.1. Deriving Theorem 2.13 from Theorem 2.9

Let us see how Theorem 2.13 can be proven using Theorem 2.9 (which we have not proven yet). We shall need two lemmas:

**Lemma 3.1.** Let  $n \in \mathbb{N}$  and  $m \in \mathbb{N}$ . Let  $b_{i,k}$  be an element of  $\mathbb{K}$  for every  $i \in \{1, 2, \dots, m\}$  and every  $k \in \{1, 2, \dots, n\}$ . Let  $d_{i,j,k}$  be an element of  $\mathbb{K}$  for every  $i \in \{1, 2, \dots, m\}$ ,  $j \in \{1, 2, \dots, m\}$  and  $k \in \{1, 2, \dots, n\}$ . Let  $G$  be the  $m \times m$ -matrix  $\left( \sum_{k=1}^n b_{i,k} d_{i,j,k} \right)_{1 \leq i \leq m, 1 \leq j \leq m}$ . Then,

$$\det G = \sum_{f: \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, n\}} \left( \prod_{i=1}^m b_{i, f(i)} \right) \det \left( \left( d_{i,j, f(i)} \right)_{1 \leq i \leq m, 1 \leq j \leq m} \right).$$

Lemma 3.1 is merely a scary way to state the multilinearity of the determinant as a function of its rows. See the Appendix for a proof.

Let us specialize Lemma 3.1 in a way that is closer to our goal:

**Lemma 3.2.** Let  $n$  be a positive integer. Let  $b_{i,k}$  be an element of  $\mathbb{K}$  for every  $i \in \{1, 2, \dots, n-1\}$  and every  $k \in \{1, 2, \dots, n\}$ . Let  $d_{i,j,k}$  be an element of  $\mathbb{K}$  for every  $i \in \{1, 2, \dots, n-1\}$ ,  $j \in \{1, 2, \dots, n-1\}$  and  $k \in \{1, 2, \dots, n\}$ . Let  $G$  be the  $(n-1) \times (n-1)$ -matrix  $\left( \sum_{k=1}^n b_{i,k} d_{i,j,k} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1}$ . Then,

$$\det G = \sum_{\substack{f: \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}; \\ f(n) = n}} \left( \prod_{i=1}^{n-1} b_{i, f(i)} \right) \det \left( \left( d_{i,j, f(i)} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right).$$

*Proof of Lemma 3.2.* Lemma 3.1 (applied to  $m = n - 1$ ) shows that

$$\det G = \sum_{f: \{1, 2, \dots, n-1\} \rightarrow \{1, 2, \dots, n\}} \left( \prod_{i=1}^{n-1} b_{i, f(i)} \right) \det \left( \left( d_{i,j, f(i)} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right).$$

The only difference between this formula and the claim of Lemma 3.2 is that the sum here is over all  $f : \{1, 2, \dots, n-1\} \rightarrow \{1, 2, \dots, n\}$ , whereas the sum in the claim of Lemma 3.2 is over all  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  satisfying  $f(n) = n$ . But this is not much of a difference: Each map  $\{1, 2, \dots, n-1\} \rightarrow \{1, 2, \dots, n\}$  is a restriction (to  $\{1, 2, \dots, n-1\}$ ) of a unique map  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  satisfying  $f(n) = n$ , and therefore the two sums are equal.  $\square$



*Proof of Theorem 2.13.* For every  $i \in \{1, 2, \dots, n-1\}$ ,  $j \in \{1, 2, \dots, n-1\}$  and  $k \in \{1, 2, \dots, n\}$ , define an element  $d_{i,j,k}$  of  $\mathbb{K}$  by

$$d_{i,j,k} = a_{i,j}a_{k,n} - a_{i,n}a_{k,j}. \tag{2}$$

For every  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  satisfying  $f(n) = n$ , we have

$$\begin{aligned} & \det \left( \left( \begin{array}{c} \underbrace{d_{i,j,f(i)}}_{=a_{i,j}a_{f(i),n} - a_{i,n}a_{f(i),j}} \\ \text{(by (2))} \end{array} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) \\ &= \det \left( \left( a_{i,j}a_{f(i),n} - a_{i,n}a_{f(i),j} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) \\ &= \begin{cases} 0, & \text{if } f \text{ is not } n\text{-potent;} \\ (\text{abut}_f A) \cdot \det A, & \text{if } f \text{ is } n\text{-potent} \end{cases} \end{aligned} \tag{3}$$

(by Theorem 2.9, applied to the matrix  $\left( a_{i,j}a_{f(i),n} - a_{i,n}a_{f(i),j} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1}$  instead of  $B$ ).

We have

$$(c_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} = BA = \left( \sum_{k=1}^n b_{i,k}a_{k,j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}$$

(by the definition of the product of two matrices). Thus,

$$c_{i,j} = \sum_{k=1}^n b_{i,k}a_{k,j} \quad \text{for every } (i,j) \in \{1, 2, \dots, n\}^2. \tag{4}$$

Now, for every  $(i,j) \in \{1, 2, \dots, n-1\}^2$ , we have

$$\begin{aligned} & a_{i,j} \underbrace{c_{i,n}}_{= \sum_{k=1}^n b_{i,k}a_{k,n}} - a_{i,n} \underbrace{c_{i,j}}_{= \sum_{k=1}^n b_{i,k}a_{k,j}} \\ & \quad \text{(by (4), applied to } n \text{ instead of } j) \quad \text{(by (4))} \\ &= a_{i,j} \sum_{k=1}^n b_{i,k}a_{k,n} - a_{i,n} \sum_{k=1}^n b_{i,k}a_{k,j} = \sum_{k=1}^n b_{i,k} \underbrace{(a_{i,j}a_{k,n} - a_{i,n}a_{k,j})}_{=d_{i,j,k}} = \sum_{k=1}^n b_{i,k}d_{i,j,k}. \\ & \quad \text{(by (2))} \end{aligned}$$

Hence,

$$G = \left( \begin{array}{c} \underbrace{a_{i,j}c_{i,n} - a_{i,n}c_{i,j}}_{= \sum_{k=1}^n b_{i,k}d_{i,j,k}} \\ \end{array} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} = \left( \sum_{k=1}^n b_{i,k}d_{i,j,k} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1}.$$

Hence, Lemma 3.2 yields

$$\begin{aligned}
 \det G &= \sum_{\substack{f:\{1,2,\dots,n\}\rightarrow\{1,2,\dots,n\}; \\ f(n)=n}} \underbrace{\left( \prod_{i=1}^{n-1} b_{i,f(i)} \right)}_{\substack{=\text{weight}_f B \\ \text{(by the definition} \\ \text{of } \text{weight}_f B)}} \underbrace{\det \left( \left( d_{i,j,f(i)} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right)}_{\substack{0, & \text{if } f \text{ is not } n\text{-potent;} \\ (\text{abut}_f A) \cdot \det A, & \text{if } f \text{ is } n\text{-potent}}} \\
 &= \sum_{\substack{f:\{1,2,\dots,n\}\rightarrow\{1,2,\dots,n\}; \\ f(n)=n}} \left( \text{weight}_f B \right) \begin{cases} 0, & \text{if } f \text{ is not } n\text{-potent;} \\ (\text{abut}_f A) \cdot \det A, & \text{if } f \text{ is } n\text{-potent} \end{cases} \\
 &= \sum_{\substack{f:\{1,2,\dots,n\}\rightarrow\{1,2,\dots,n\}; \\ f(n)=n; \\ f \text{ is } n\text{-potent}}} \left( \text{weight}_f B \right) (\text{abut}_f A) \cdot \det A \\
 &= \left( \sum_{\substack{f:\{1,2,\dots,n\}\rightarrow\{1,2,\dots,n\}; \\ f(n)=n; \\ f \text{ is } n\text{-potent}}} \left( \text{weight}_f B \right) (\text{abut}_f A) \right) \cdot \det A.
 \end{aligned}$$

□

### 3.2. Deriving Theorem 2.12 from Theorem 2.13

Now let us see why Theorem 2.13 generalizes the matrix-tree theorem.

*Proof of Theorem 2.12.* WLOG assume that  $n \geq 2$  (since the case  $n = 1$  is easy to check by hand). Define an  $n \times n$ -matrix  $A$  by  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ , where

$$a_{i,j} = \delta_{i,j} + \delta_{j,n} (1 - \delta_{i,n}).$$

(This scary formula hides a simple idea: this is the matrix whose entries on the diagonal and in its last column are 1, and all other entries are 0. Thus,

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 1 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 1 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & 1 & \cdots & 0 & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 1 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix}.$$

) Note that every  $(i, j) \in \{1, 2, \dots, n-1\}^2$  satisfies

$$a_{i,j} = \delta_{i,j} + \underbrace{\delta_{j,n}}_{\substack{=0 \\ \text{(since } j \neq n \\ \text{(since } j \in \{1, 2, \dots, n-1\})})}} (1 - \delta_{i,n}) = \delta_{i,j}. \quad (5)$$

Also, every  $i \in \{1, 2, \dots, n-1\}$  satisfies

$$\begin{aligned} a_{i,n} &= \underbrace{\delta_{i,n}}_{\substack{=0 \\ \text{(since } i \neq n)}} + \underbrace{\delta_{n,n}}_{=1} \left( 1 - \underbrace{\delta_{i,n}}_{\substack{=0 \\ \text{(since } i \neq n)}} \right) && \text{(by the definition of } a_{i,n}) \\ &= 0 + 1(1 - 0) = 1. \end{aligned} \quad (6)$$

Also, let  $B$  be the  $n \times n$ -matrix  $(W(i, j))_{1 \leq i \leq n, 1 \leq j \leq n}$ . Write the  $n \times n$ -matrix  $BA$  in the form  $BA = (c_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ . Then, it is easy to see that every  $(i, j) \in \{1, 2, \dots, n\}^2$  satisfies

$$c_{i,j} = W(i, j) + \delta_{j,n} (d^+(i) - W(i, n)) \quad (7)$$

6.

---

<sup>6</sup>Proof of (7): For every  $i \in \{1, 2, \dots, n\}$ , we have

$$\begin{aligned} d^+(i) &= \sum_{j=1}^n W(i, j) && \text{(by the definition of } d^+(i)) \\ &= \sum_{j=1}^{n-1} W(i, j) + W(i, n) = \sum_{k=1}^{n-1} W(i, k) + W(i, n) \end{aligned}$$

(here, we renamed the summation index  $j$  as  $k$ ) and thus

$$\sum_{k=1}^{n-1} W(i, k) = d^+(i) - W(i, n). \quad (8)$$

But

$$(c_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} = BA = \left( \sum_{k=1}^n W(i, k) a_{k,j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}$$

(by the definition of the product of two matrices, since  $B = (W(i, j))_{1 \leq i \leq n, 1 \leq j \leq n}$  and  $A =$

---

Thus, for every  $(i, j) \in \{1, 2, \dots, n-1\}^2$ , we have

$$\begin{aligned}
 & \underbrace{a_{i,j}}_{=\delta_{i,j} \text{ (by (5))}} = \underbrace{c_{i,n}}_{=\underbrace{W(i,n) + \delta_{n,n}(d^+(i) - W(i,n))}_{\text{(by (7), applied to } j \text{ instead of } n)}} - \underbrace{a_{i,n}}_{=\underbrace{1}_{\text{(by (6))}}} = \underbrace{c_{i,j}}_{=\underbrace{W(i,j) + \delta_{j,n}(d^+(i) - W(i,n))}_{\text{(by (7))}}} \\
 & = \delta_{i,j} \left( W(i,n) + \underbrace{\delta_{n,n}}_{=1} (d^+(i) - W(i,n)) \right) - \left( W(i,j) + \underbrace{\delta_{j,n}}_{=\underbrace{0}_{\text{(since } j < n)}} (d^+(i) - W(i,n)) \right) \\
 & = \delta_{i,j} \underbrace{(W(i,n) + (d^+(i) - W(i,n)))}_{=d^+(i)} - W(i,j) = \delta_{i,j} d^+(i) - W(i,j).
 \end{aligned}$$

Hence,

$$(a_{i,j}c_{i,n} - a_{i,n}c_{i,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} = (\delta_{i,j}d^+(i) - W(i,j))_{1 \leq i \leq n-1, 1 \leq j \leq n-1} = L.$$

In other words,  $L$  is the matrix  $(a_{i,j}c_{i,n} - a_{i,n}c_{i,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \in \mathbb{K}^{(n-1) \times (n-1)}$ .

$(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ . Hence, every  $(i, j) \in \{1, 2, \dots, n\}^2$  satisfies

$$\begin{aligned}
 c_{i,j} &= \sum_{k=1}^n W(i,k) \underbrace{a_{k,j}}_{=\underbrace{\delta_{k,j} + \delta_{j,n}(1 - \delta_{k,n})}_{\text{(by the definition of } a_{k,j})}} \\
 &= \sum_{k=1}^n W(i,k) (\delta_{k,j} + \delta_{j,n}(1 - \delta_{k,n})) \\
 &= \underbrace{\sum_{k=1}^n W(i,k) \delta_{k,j}}_{=\underbrace{W(i,j)}_{\text{(because the factor } \delta_{k,j} \text{ in the sum kills every addend except the one for } k=j)}} + \delta_{j,n} \underbrace{\sum_{k=1}^n W(i,k) (1 - \delta_{k,n})}_{=\sum_{k=1}^{n-1} W(i,k)(1 - \delta_{k,n}) + W(i,n)(1 - \delta_{n,n})} \\
 &= W(i,j) + \delta_{j,n} \left( \sum_{k=1}^{n-1} W(i,k) \left( 1 - \underbrace{\delta_{k,n}}_{=\underbrace{0}_{\text{(since } k < n)}} \right) + W(i,n) \underbrace{(1 - \delta_{n,n})}_{=\underbrace{0}_{\text{(since } \delta_{n,n}=1)}} \right) \\
 &= W(i,j) + \delta_{j,n} \left( \sum_{k=1}^{n-1} W(i,k) \underbrace{(1 - 0)}_{=1} + \underbrace{W(i,n) 0}_{=0} \right) \\
 &= W(i,j) + \delta_{j,n} \underbrace{\sum_{k=1}^{n-1} W(i,k)}_{=\underbrace{d^+(i) - W(i,n)}_{\text{(by (8))}}} = W(i,j) + \delta_{j,n} (d^+(i) - W(i,n)),
 \end{aligned}$$

and thus (7) is proven.

Thus, Theorem 2.13 (applied to  $G = L$ ) yields

$$\det L = \left( \sum_{\substack{f: \{1,2,\dots,n\} \rightarrow \{1,2,\dots,n\}; \\ f(n)=n; \\ f \text{ is } n\text{-potent}}} \underbrace{\left( \text{weight}_f B \right)}_{= \prod_{i=1}^{n-1} W(i, f(i))} \underbrace{\left( \text{abut}_f A \right)}_{=1} \right) \cdot \underbrace{\det A}_{=1}$$

$$= \sum_{\substack{f: \{1,2,\dots,n\} \rightarrow \{1,2,\dots,n\}; \\ f(n)=n; \\ f \text{ is } n\text{-potent}}} \prod_{i=1}^{n-1} W(i, f(i)).$$

This proves Theorem 2.12. □

### 3.3. Some combinatorial lemmas

We still owe the reader a proof of Theorem 2.9. We prepare by proving some properties of maps  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ .

**Proposition 3.3.** Let  $n \in \mathbb{N}$ . Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map. Let  $i \in \{1, 2, \dots, n\}$ . Then,

$$f^k(i) \in \{f^s(i) \mid s \in \{0, 1, \dots, n-1\}\} \quad \text{for every } k \in \mathbb{N}.$$

Proposition 3.3 is a classical fact; we give the proof in the Appendix below.

The following three results can be easily derived from Proposition 3.3; we shall give more detailed proofs in the Appendix:

**Proposition 3.4.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map such that  $f(n) = n$ . Let  $i \in \{1, 2, \dots, n\}$ . Then,  $f^{n-1}(i) = n$  if and only if there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$ .

**Proposition 3.5.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map such that  $f(n) = n$ . Then, the map  $f$  is  $n$ -potent if and only if  $f^{n-1}(\{1, 2, \dots, n\}) = \{n\}$ .

**Corollary 3.6.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map such that  $f(n) = n$ . Let  $i \in \{1, 2, \dots, n\}$ . Then,  $\delta_{f^{n-1}(i), n} = \delta_{f^n(i), n}$ .

One consequence of Proposition 3.5 is the following: If  $n$  is a positive integer, and if  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  is a map such that  $f(n) = n$ , then we can check in finite time whether the map  $f$  is  $n$ -potent (because we can check in finite time whether  $f^{n-1}(\{1, 2, \dots, n\}) = \{n\}$ ). Thus, for any given positive integer  $n$ , it is possible to enumerate all  $n$ -potent maps  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ .

Next, we shall show a property of  $n$ -potent maps:

**Lemma 3.7.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map such that  $f(n) = n$ . Assume that  $f$  is  $n$ -potent.

Let  $\sigma \in S_n$  be a permutation such that  $\sigma \neq \text{id}$ . Then, there exists some  $i \in \{1, 2, \dots, n\}$  such that  $\sigma(i) \notin \{i, f(i)\}$ .

*Proof of Lemma 3.7.* Assume the contrary. Thus,  $\sigma(i) \in \{i, f(i)\}$  for every  $i \in \{1, 2, \dots, n\}$ .

We have  $\sigma \neq \text{id}$ . Hence, there exists some  $h \in \{1, 2, \dots, n\}$  such that  $\sigma(h) \neq h$ . Fix such a  $h$ . We shall prove that

$$\sigma^j(h) = f^j(h) \quad \text{for every } j \in \mathbb{N}. \tag{9}$$

Indeed, we shall prove this by induction over  $j$ . The induction base (the case  $j = 0$ ) is obvious. For the induction step, fix  $J \in \mathbb{N}$ , and assume that  $\sigma^J(h) = f^J(h)$ . We need to prove that  $\sigma^{J+1}(h) = f^{J+1}(h)$ .

We have assumed that  $\sigma(i) \in \{i, f(i)\}$  for every  $i \in \{1, 2, \dots, n\}$ . Applying this to  $i = \sigma^J(h)$ , we obtain  $\sigma(\sigma^J(h)) \in \{\sigma^J(h), f(\sigma^J(h))\}$ . In other words,  $\sigma^{J+1}(h) \in \{\sigma^J(h), f(\sigma^J(h))\}$ . Thus, either  $\sigma^{J+1}(h) = \sigma^J(h)$  or  $\sigma^{J+1}(h) = f(\sigma^J(h))$ . Since  $\sigma^{J+1}(h) = \sigma^J(h)$  is impossible (because in light of the invertibility of  $\sigma$ , this would yield  $\sigma(h) = h$ , which contradicts  $\sigma(h) \neq h$ ), we thus must have  $\sigma^{J+1}(h) =$

$$f(\sigma^J(h)). \text{ Hence, } \sigma^{J+1}(h) = f\left(\underbrace{\sigma^J(h)}_{=f^J(h)}\right) = f(f^J(h)) = f^{J+1}(h). \text{ This completes}$$

the induction step.

Thus, (9) is proven.

But  $f$  is  $n$ -potent. Hence, there exists some  $k \in \mathbb{N}$  such that  $f^k(h) = n$ . Consider this  $k$ . Applying (9) to  $j = k$ , we obtain  $\sigma^k(h) = f^k(h) = n$ .

But applying (9) to  $j = k + 1$ , we obtain  $\sigma^{k+1}(h) = f^{k+1}(h) = f\left(\underbrace{f^k(h)}_{=n}\right) = f(n) = n$ . Hence,  $n = \sigma^{k+1}(h) = \sigma^k(\sigma(h))$ , so that  $\sigma^k(\sigma(h)) = n = \sigma^k(h)$ . Since  $\sigma^k$  is invertible, this entails  $\sigma(h) = h$ , which contradicts  $\sigma(h) \neq h$ . This contradiction proves that our assumption was wrong. Thus, Lemma 3.7 is proven.  $\square$

### 3.4. The matrix $Z_f$ and its determinant

Next, we assign a matrix  $Z_f$  to every such  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ :

**Definition 3.8.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map. Then, we define an  $n \times n$ -matrix  $Z_f \in \mathbb{K}^{n \times n}$  by

$$Z_f = \left( \delta_{i,j} - (1 - \delta_{i,n}) \delta_{f(i),j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}.$$

**Example 3.9.** For this example, set  $n = 4$ , and define a map  $f : \{1, 2, 3, 4\} \rightarrow \{1, 2, 3, 4\}$  by  $(f(1), f(2), f(3), f(4)) = (2, 4, 1, 4)$ . Then,

$$Z_f = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Now, we claim the following:

**Proposition 3.10.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map such that  $f(n) = n$ . Let  $v_f$  be the column vector  $\left(1 - \delta_{f^{n-1}(i), n}\right)_{1 \leq i \leq n, 1 \leq j \leq 1} \in \mathbb{K}^{n \times 1}$ . Then,  $Z_f v_f = 0_{n \times 1}$ .

(Recall that  $0_{n \times 1}$  denotes the  $n \times 1$  zero matrix, i.e., the column vector with  $n$  entries whose all entries are 0.)

*Proof of Proposition 3.10.* We shall prove that

$$\sum_{k=1}^n \left( \delta_{i,k} - (1 - \delta_{i,n}) \delta_{f(i),k} \right) \left( 1 - \delta_{f^{n-1}(k), n} \right) = 0 \quad (10)$$

for every  $i \in \{1, 2, \dots, n\}$ .

*Proof of (10):* Let  $i \in \{1, 2, \dots, n\}$ . Corollary 3.6 yields  $\delta_{f^{n-1}(i), n} = \delta_{f^n(i), n}$ .

On the other hand,  $f(n) = n$ . Thus, it is straightforward to see (by induction over  $h$ ) that  $f^h(n) = n$  for every  $h \in \mathbb{N}$ . Applying this to  $h = n$ , we obtain  $f^n(n) = n$ .

Now,

$$\begin{aligned}
 & \sum_{k=1}^n \left( \delta_{i,k} - (1 - \delta_{i,n}) \delta_{f(i),k} \right) \left( 1 - \delta_{f^{n-1}(k),n} \right) \\
 &= \underbrace{\sum_{k=1}^n \delta_{i,k} \left( 1 - \delta_{f^{n-1}(k),n} \right)}_{=1 - \delta_{f^{n-1}(i),n}} - \underbrace{\sum_{k=1}^n (1 - \delta_{i,n}) \delta_{f(i),k} \left( 1 - \delta_{f^{n-1}(k),n} \right)}_{=(1 - \delta_{i,n}) \left( 1 - \delta_{f^{n-1}(f(i)),n} \right)} \\
 & \quad \text{(because the factor } \delta_{i,k} \text{ in the sum kills every addend except the one for } k=i) \quad \text{(because the factor } \delta_{f(i),k} \text{ in the sum kills every addend except the one for } k=f(i)) \\
 &= \left( 1 - \underbrace{\delta_{f^{n-1}(i),n}}_{=\delta_{f^n(i),n}} \right) - (1 - \delta_{i,n}) \left( 1 - \underbrace{\delta_{f^{n-1}(f(i)),n}}_{=\delta_{f^n(i),n}} \right) \\
 &= \left( 1 - \delta_{f^n(i),n} \right) - (1 - \delta_{i,n}) \left( 1 - \delta_{f^n(i),n} \right) \\
 &= \underbrace{\left( 1 - (1 - \delta_{i,n}) \right)}_{=\delta_{i,n}} \left( 1 - \delta_{f^n(i),n} \right) = \delta_{i,n} \left( 1 - \delta_{f^n(i),n} \right) \\
 &= \begin{cases} 0, & \text{if } i \neq n; \\ 1 - \delta_{f^n(n),n}, & \text{if } i = n \end{cases} = \begin{cases} 0, & \text{if } i \neq n; \\ 0, & \text{if } i = n \end{cases} \\
 & \quad \left( \text{since } f^n(n) = n \text{ and thus } \delta_{f^n(n),n} = \delta_{n,n} = 1 \text{ and hence } 1 - \delta_{f^n(n),n} = 0 \right) \\
 &= 0.
 \end{aligned}$$

This proves (10).

Recall now that

$$Z_f = \left( \delta_{i,j} - (1 - \delta_{i,n}) \delta_{f(i),j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}$$

and  $v_f = \left( 1 - \delta_{f^{n-1}(i),n} \right)_{1 \leq i \leq n, 1 \leq j \leq 1}$ . Hence, the definition of the product of two matrices yields

$$\begin{aligned}
 Z_f v_f &= \left( \underbrace{\sum_{k=1}^n \left( \delta_{i,k} - (1 - \delta_{i,n}) \delta_{f(i),k} \right) \left( 1 - \delta_{f^{n-1}(k),n} \right)}_{\substack{=0 \\ \text{(by (10))}}} \right)_{1 \leq i \leq n, 1 \leq j \leq 1} \\
 &= (0)_{1 \leq i \leq n, 1 \leq j \leq 1} = 0_{n \times 1}.
 \end{aligned}$$

This proves Proposition 3.10. □



Now, we recall the following well-known properties of determinants<sup>7</sup>:

**Lemma 3.11.** Let  $n \in \mathbb{N}$ . Let  $A$  be an  $n \times n$ -matrix. Let  $v$  be a column vector with  $n$  entries. If  $Av = 0_{n \times 1}$ , then  $\det A \cdot v = 0_{n \times 1}$ .

**Lemma 3.12.** Let  $n$  be a positive integer. Let  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$  be an  $n \times n$ -matrix. Assume that

$$a_{i,n} = 0 \quad \text{for every } i \in \{1, 2, \dots, n-1\}. \quad (11)$$

Then,  $\det A = a_{n,n} \cdot \det \left( (a_{i,j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right)$ .

Now, we can prove the crucial property of the matrix  $Z_f$ :

**Proposition 3.13.** Let  $n$  be a positive integer. Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be a map satisfying  $f(n) = n$ .

- (a) If  $f$  is  $n$ -potent, then  $\det(Z_f) = 1$ .
- (b) If  $f$  is not  $n$ -potent, then  $\det(Z_f) = 0$ .

*Proof of Proposition 3.13.* Write the matrix  $Z_f$  in the form  $(z_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ . Thus,

$$(z_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} = Z_f = \left( \delta_{i,j} - (1 - \delta_{i,n}) \delta_{f(i),j} \right)_{1 \leq i \leq n, 1 \leq j \leq n}.$$

Hence, every  $(i, j) \in \{1, 2, \dots, n\}^2$  satisfies

$$z_{i,j} = \delta_{i,j} - \underbrace{(1 - \delta_{i,n})}_{\delta_{f(i),j}} \delta_{f(i),j} = \delta_{i,j} - \begin{cases} 1, & \text{if } i < n; \\ 0, & \text{if } i = n \end{cases} \delta_{f(i),j} \quad (12)$$

$$= \begin{cases} 1, & \text{if } i < n; \\ 0, & \text{if } i = n \end{cases}$$

$$= \delta_{i,j} - \begin{cases} \delta_{f(i),j'} & \text{if } i < n; \\ 0, & \text{if } i = n \end{cases} = \begin{cases} \delta_{i,j} - \delta_{f(i),j'} & \text{if } i < n; \\ \delta_{i,j'} & \text{if } i = n \end{cases}. \quad (13)$$

(a) Assume that  $f$  is  $n$ -potent.

Let  $\sigma \in S_n$  be a permutation such that  $\sigma \neq \text{id}$ . Then, there exists some  $i \in \{1, 2, \dots, n\}$  such that  $\sigma(i) \notin \{i, f(i)\}$  (by Lemma 3.7). Hence, there exists some  $i \in \{1, 2, \dots, n\}$  such that  $z_{i,\sigma(i)} = 0$ <sup>8</sup>. Hence, the product  $\prod_{i=1}^n z_{i,\sigma(i)}$  has at least one zero factor, and thus equals 0.

<sup>7</sup>For the sake of completeness: Lemma 3.11 is [Grinbe15, Corollary 6.102]; Lemma 3.12 is [Grinbe15, Corollary 6.45].

<sup>8</sup>*Proof.* We have just shown that there exists some  $i \in \{1, 2, \dots, n\}$  such that  $\sigma(i) \notin \{i, f(i)\}$ . Consider this  $i$ . We have  $\sigma(i) \notin \{i, f(i)\}$ , thus  $\sigma(i) \neq i$ , and thus  $\delta_{i,\sigma(i)} = 0$ . Also,  $\sigma(i) \notin$

Now, forget that we fixed  $\sigma$ . We thus have shown that

$$\prod_{i=1}^n z_{i,\sigma(i)} = 0 \quad \text{for every } \sigma \in S_n \text{ such that } \sigma \neq \text{id}. \quad (14)$$

On the other hand, it is easy to see that

$$\prod_{i=1}^n z_{i,i} = 1. \quad (15)$$

9

Now, the definition of  $\det(Z_f)$  yields

$$\begin{aligned} \det(Z_f) &= \sum_{\sigma \in S_n} (-1)^\sigma \prod_{i=1}^n z_{i,\sigma(i)} \quad \left( \text{since } Z_f = (z_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \right) \\ &= \underbrace{(-1)^{\text{id}}}_{=1} \prod_{i=1}^n \underbrace{z_{i,\text{id}(i)}}_{=z_{i,i}} + \sum_{\substack{\sigma \in S_n; \\ \sigma \neq \text{id}}} (-1)^\sigma \underbrace{\prod_{i=1}^n z_{i,\sigma(i)}}_{=0} \\ &\hspace{15em} \text{(by (14))} \\ &= \prod_{i=1}^n z_{i,i} + \underbrace{\sum_{\substack{\sigma \in S_n; \\ \sigma \neq \text{id}}} (-1)^\sigma 0}_{=0} = \prod_{i=1}^n z_{i,i} = 1 \quad \text{(by (15)).} \end{aligned}$$

This proves Proposition 3.13 (a).

(b) Assume that  $f$  is not  $n$ -potent. Then, there exists some  $i \in \{1, 2, \dots, n\}$  such that  $f^{n-1}(i) \neq n$ <sup>10</sup>. Fix such an  $i$ , and denote it by  $u$ . Thus,  $u \in \{1, 2, \dots, n\}$  is such that  $f^{n-1}(u) \neq n$ .

---

$\{i, f(i)\}$ , thus  $\sigma(i) \neq f(i)$ , and thus  $\delta_{f(i),\sigma(i)} = 0$ . Now, (12) (applied to  $(i, \sigma(i))$  instead of  $(i, j)$ ) yields

$$z_{i,\sigma(i)} = \underbrace{\delta_{i,\sigma(i)}}_{=0} - \begin{cases} 1, & \text{if } i < n; \\ 0, & \text{if } i = n \end{cases} \underbrace{\delta_{f(i),\sigma(i)}}_{=0} = 0 - 0 = 0,$$

qed.

<sup>9</sup>Proof of (15): To prove this, it is sufficient to show that  $z_{i,i} = 1$  for every  $i \in \{1, 2, \dots, n\}$ . This is obvious when  $i = n$  (using the formula (13)), so we only need to consider the case when  $i < n$ . In this case, (13) (applied to  $(i, i)$  instead of  $(i, j)$ ) shows that  $z_{i,i} = \underbrace{\delta_{i,i}}_{=1} - \delta_{f(i),i} = 1 - \delta_{f(i),i}$ .

Hence, in order to prove that  $z_{i,i} = 1$ , we need to show that  $\delta_{f(i),i} = 0$ . In other words, we need to prove that  $f(i) \neq i$ .

Indeed, assume the contrary. Thus,  $f(i) = i$ . Hence, by induction over  $k$ , we can easily see that  $f^k(i) = i$  for every  $k \in \mathbb{N}$ . Hence, for every  $k \in \mathbb{N}$ , we have  $f^k(i) = i \neq n$ . This contradicts the fact that there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (since  $f$  is  $n$ -potent). This contradiction proves that our assumption was wrong. Hence, (15) is proven.

<sup>10</sup>Proof. Assume the contrary. Thus, for every  $i \in \{1, 2, \dots, n\}$ , we have  $f^{n-1}(i) = n$ . Hence, for every  $i \in \{1, 2, \dots, n\}$ , there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (according to the  $\implies$  direction of Proposition 3.4). In other words, the map  $f$  is  $n$ -potent. This contradicts the fact that  $f$  is not  $n$ -potent. This contradiction shows that our assumption was wrong, qed.

Define the vector  $v_f$  as in Proposition 3.10. Proposition 3.10 yields  $Z_f v_f = 0_{n \times 1}$ . Lemma 3.11 (applied to  $Z_f$  and  $v_f$  instead of  $A$  and  $v$ ) thus yields  $\det(Z_f) \cdot v_f = 0_{n \times 1}$ . Thus,

$$\begin{aligned} (0)_{1 \leq i \leq n, 1 \leq j \leq 1} &= 0_{n \times 1} = \det(Z_f) \cdot \underbrace{v_f}_{=(1 - \delta_{f^{n-1}(i),n})_{1 \leq i \leq n, 1 \leq j \leq 1}} \\ &= \det(Z_f) \cdot (1 - \delta_{f^{n-1}(i),n})_{1 \leq i \leq n, 1 \leq j \leq 1} \\ &= \left( \det(Z_f) \cdot (1 - \delta_{f^{n-1}(i),n}) \right)_{1 \leq i \leq n, 1 \leq j \leq 1}. \end{aligned}$$

In other words,  $0 = \det(Z_f) \cdot (1 - \delta_{f^{n-1}(i),n})$  for each  $i \in \{1, 2, \dots, n\}$ . Applying this to  $i = u$ , we obtain

$$0 = \det(Z_f) \cdot \left( 1 - \underbrace{\delta_{f^{n-1}(u),n}}_{\substack{=0 \\ \text{(since } f^{n-1}(u) \neq n)}}} \right) = \det(Z_f) \cdot 1 = \det(Z_f).$$

This proves Proposition 3.13 (b). □

### 3.5. Proof of Theorem 2.9

Let us finally recall a particularly basic property of determinants:

**Lemma 3.14.** Let  $m \in \mathbb{N}$ . Let  $A = (a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq m} \in \mathbb{K}^{m \times m}$  be an  $m \times m$ -matrix. Let  $b_1, b_2, \dots, b_m$  be  $m$  elements of  $\mathbb{K}$ . Then,

$$\det\left((b_i a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq m}\right) = \left(\prod_{i=1}^m b_i\right) \det A.$$

(Again, see the Appendix for the proof of this lemma.)

We can now finally prove Theorem 2.9:

*Proof of Theorem 2.9.* The identities we want to prove (both for part (a) and for part (b)) are polynomial identities in the entries of  $A$ . Thus, we can WLOG assume that all these entries are invertible.<sup>11</sup> In other words, we can assume that  $a_{i,j}$  is invertible for each  $(i, j) \in \{1, 2, \dots, n\}^2$ . Assume this.

<sup>11</sup>Here is a more detailed justification for this “WLOG”:

Let us restrict ourselves to Theorem 2.9 (b). (The argument for Theorem 2.9 (a) is analogous.)

Assume that Theorem 2.9 (b) is proven in the case when all entries of  $A$  are invertible. We now must show that Theorem 2.9 (b) always holds.

Let  $C$  be the  $(n - 1) \times (n - 1)$ -matrix

$$\left( \begin{array}{cc} a_{i,j} & a_{f(i),j} \\ a_{i,n} & a_{f(i),n} \end{array} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \in \mathbb{K}^{(n-1) \times (n-1)}.$$

Let  $n$  be a positive integer such that  $n \geq 2$ . Let  $f : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be an  $n$ -potent map. Then, Theorem 2.9 **(b)** claims that

$$\det B = (\text{abut}_f A) \cdot \det A \tag{16}$$

for every  $n \times n$ -matrix  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$ , where  $B$  is as defined in Theorem 2.9. The equality (16) rewrites as

$$\begin{aligned} & \sum_{\sigma \in S_{n-1}} \prod_{i=1}^{n-1} (a_{i,\sigma(i)} a_{f(i),n} - a_{i,n} a_{f(i),\sigma(i)}) \\ &= \left( a_{n,n}^{|f^{-1}(n)|-2} \prod_{\substack{i \in \{1,2,\dots,n-1\}; \\ f(i) \neq n}} a_{f(i),n} \right) \cdot \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)} \end{aligned} \tag{17}$$

(because we have

$$\begin{aligned} \det \underbrace{B}_{=(a_{i,j} a_{f(i),n} - a_{i,n} a_{f(i),j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1}} &= \det \left( (a_{i,j} a_{f(i),n} - a_{i,n} a_{f(i),j})_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) \\ &= \sum_{\sigma \in S_{n-1}} \prod_{i=1}^{n-1} (a_{i,\sigma(i)} a_{f(i),n} - a_{i,n} a_{f(i),\sigma(i)}) \end{aligned}$$

and  $\text{abut}_f A = a_{n,n}^{|f^{-1}(n)|-2} \prod_{\substack{i \in \{1,2,\dots,n-1\}; \\ f(i) \neq n}} a_{f(i),n}$  and  $\det A = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)}$ ). Thus, Theorem 2.9

**(b)** (for our given  $n$  and  $f$ ) is equivalent to the claim that (17) holds for every  $n \times n$ -matrix  $(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$ .

Now, let  $\mathbb{P}$  be the polynomial ring  $\mathbb{Z} [X_{i,j} \mid (i,j) \in \{1, 2, \dots, n\}^2]$  in the  $n^2$  indeterminates  $X_{i,j}$  for  $(i,j) \in \{1, 2, \dots, n\}^2$ . Let  $\mathbb{F}$  be the quotient field of  $\mathbb{P}$ ; this is the field  $\mathbb{Q} (X_{i,j} \mid (i,j) \in \{1, 2, \dots, n\}^2)$  of rational functions in the same indeterminates (but over  $\mathbb{Q}$ ).

Let  $A_X$  be the  $n \times n$ -matrix  $(X_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{P}^{n \times n}$ . If we regard  $A_X$  as a matrix in  $\mathbb{F}^{n \times n}$ , then all entries of  $A_X$  are invertible (because they are nonzero elements of the field  $\mathbb{F}$ ). Hence, Theorem 2.9 **(b)** can be applied to  $\mathbb{F}$ ,  $A_X$ ,  $X_{i,j}$  and  $B_X$  instead of  $\mathbb{K}$ ,  $A$ ,  $a_{i,j}$  and  $B$  (because we have assumed that Theorem 2.9 **(b)** is proven in the case when all entries of  $A$  are invertible). As we know, this means that (17) holds for  $a_{i,j} = X_{i,j}$ . In other words, we have

$$\begin{aligned} & \sum_{\sigma \in S_{n-1}} \prod_{i=1}^{n-1} (X_{i,\sigma(i)} X_{f(i),n} - X_{i,n} X_{f(i),\sigma(i)}) \\ &= \left( X_{n,n}^{|f^{-1}(n)|-2} \prod_{\substack{i \in \{1,2,\dots,n-1\}; \\ f(i) \neq n}} X_{f(i),n} \right) \cdot \sum_{\sigma \in S_n} \prod_{i=1}^n X_{i,\sigma(i)}. \end{aligned} \tag{18}$$

Thus, Lemma 3.14 (applied to  $n - 1$ ,  $C$ ,  $\frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}}$  and  $a_{i,n}a_{f(i),n}$  instead of  $m$ ,  $A$ ,  $a_{i,j}$  and  $b_i$ ) yields

$$\begin{aligned} & \det \left( \left( a_{i,n}a_{f(i),n} \left( \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}} \right) \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) \\ &= \left( \prod_{i=1}^{n-1} (a_{i,n}a_{f(i),n}) \right) \det C. \end{aligned}$$

Comparing this with

$$\begin{aligned} & \det \left( \left( \underbrace{a_{i,n}a_{f(i),n} \left( \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}} \right)}_{=a_{i,j}a_{f(i),n} - a_{i,n}a_{f(i),j}} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right) \\ &= \det \left( \underbrace{(a_{i,j}a_{f(i),n} - a_{i,n}a_{f(i),j})}_{=B} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} = \det B, \end{aligned}$$

we find

$$\det B = \left( \prod_{i=1}^{n-1} (a_{i,n}a_{f(i),n}) \right) \det C. \tag{19}$$

It remains to compute  $\det C$ .

For every  $(i, j) \in \{1, 2, \dots, n\}^2$ , define an element  $d_{i,j} \in \mathbb{K}$  by

$$d_{i,j} = \begin{cases} \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}}, & \text{if } i < n; \\ \frac{a_{i,j}}{a_{i,n}}, & \text{if } i = n \end{cases}.$$

---

Now, let  $(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$  be an  $n \times n$ -matrix. The equality (18) is an identity between polynomials in the polynomial ring  $\mathbb{P}$ . Thus, we can substitute  $a_{i,j}$  for each  $X_{i,j}$  in this equality. As a result, we obtain the equality (17).

Thus we have shown that (17) holds for every  $n \times n$ -matrix  $(a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$ . As we have already explained, this is just a restatement of Theorem 2.9 (b); hence, Theorem 2.9 (b) is proven in full generality.

(The justification above is a typical use of the “method of universal identities”. See [Conrad09] for examples of similar justifications, albeit used in different settings.)

---

For every  $i \in \{1, 2, \dots, n-1\}$ , the definition of  $d_{i,n}$  yields

$$d_{i,n} = \begin{cases} \frac{a_{i,n}}{a_{i,n}} - \frac{a_{f(i),n}}{a_{f(i),n}}, & \text{if } i < n; \\ \frac{a_{i,n}}{a_{i,n}}, & \text{if } i = n \end{cases} = \underbrace{\frac{a_{i,n}}{a_{i,n}}}_{=1} - \underbrace{\frac{a_{f(i),n}}{a_{f(i),n}}}_{=1} \quad (\text{since } i < n)$$

$$= 1 - 1 = 0.$$

Moreover, the definition of  $d_{n,n}$  yields

$$d_{n,n} = \begin{cases} \frac{a_{n,n}}{a_{n,n}} - \frac{a_{f(n),n}}{a_{f(n),n}}, & \text{if } n < n; \\ \frac{a_{n,n}}{a_{n,n}}, & \text{if } n = n \end{cases} = \frac{a_{n,n}}{a_{n,n}} \quad (\text{since } n = n)$$

$$= 1.$$

Finally, every  $i \in \{1, 2, \dots, n-1\}$  and  $j \in \{1, 2, \dots, n\}$  satisfy

$$d_{i,j} = \begin{cases} \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}}, & \text{if } i < n; \\ \frac{a_{i,j}}{a_{i,n}}, & \text{if } i = n \end{cases} = \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}} \quad (20)$$

(since  $i < n$ ).

Now, let  $D$  be the  $n \times n$ -matrix

$$(d_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}.$$

Recall that  $d_{i,n} = 0$  for every  $i \in \{1, 2, \dots, n-1\}$ . Hence, Lemma 3.12 (applied to  $D$  and  $d_{i,j}$  instead of  $A$  and  $a_{i,j}$ ) shows that

$$\det D = \underbrace{d_{n,n}}_{=1} \det \left( \begin{pmatrix} \underbrace{d_{i,j}}_{\substack{= \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}} \\ \text{(by (20))}}} \\ a_{i,n} & a_{f(i),n} \end{pmatrix}_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right)$$

$$= \det \left( \underbrace{\begin{pmatrix} \left( \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}} \right)_{1 \leq i \leq n-1, 1 \leq j \leq n-1} \right)}_{=C} \right) = \det C.$$

Hence, (19) becomes

$$\begin{aligned} \det B &= \left( \prod_{i=1}^{n-1} (a_{i,n} a_{f(i),n}) \right) \underbrace{\det C}_{=\det D} \\ &= \left( \prod_{i=1}^{n-1} (a_{i,n} a_{f(i),n}) \right) \det D. \end{aligned} \quad (21)$$

Hence, we only need to compute  $\det D$ . How do we do this?

Let  $E$  be the  $n \times n$ -matrix  $\left( \begin{array}{c} a_{i,j} \\ a_{i,n} \end{array} \right)_{1 \leq i \leq n, 1 \leq j \leq n} \in \mathbb{K}^{n \times n}$ .

Recall that  $A = (a_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$ . Lemma 3.14 (applied to  $m = n$  and  $b_i = \frac{1}{a_{i,n}}$ ) thus yields

$$\det \left( \left( \begin{array}{c} 1 \\ a_{i,n} \end{array} a_{i,j} \right)_{1 \leq i \leq n, 1 \leq j \leq n} \right) = \left( \prod_{i=1}^n \frac{1}{a_{i,n}} \right) \det A.$$

Compared with

$$\det \left( \left( \begin{array}{c} \left( \begin{array}{c} 1 \\ a_{i,n} \end{array} a_{i,j} \right) \\ \underbrace{a_{i,j}}_{= \frac{a_{i,j}}{a_{i,n}}} \end{array} \right)_{1 \leq i \leq n, 1 \leq j \leq n} \right) = \det \left( \underbrace{\left( \begin{array}{c} a_{i,j} \\ a_{i,n} \end{array} \right)_{1 \leq i \leq n, 1 \leq j \leq n}}_{=E} \right) = \det E,$$

this yields

$$\det E = \left( \prod_{i=1}^n \frac{1}{a_{i,n}} \right) \det A. \quad (22)$$

On the other hand, recall that we have defined an  $n \times n$ -matrix  $Z_f$  in Definition 3.8. We now claim that

$$D = Z_f E. \quad (23)$$

*Proof of (23):* We have  $Z_f = (\delta_{i,j} - (1 - \delta_{i,n}) \delta_{f(i),j})_{1 \leq i \leq n, 1 \leq j \leq n}$  and

$E = \left( \begin{array}{c} a_{i,j} \\ a_{i,n} \end{array} \right)_{1 \leq i \leq n, 1 \leq j \leq n}$ . Thus, the definition of the product of two matrices yields

$$Z_f E = \left( \sum_{k=1}^n (\delta_{i,k} - (1 - \delta_{i,n}) \delta_{f(i),k}) \frac{a_{k,j}}{a_{k,n}} \right)_{1 \leq i \leq n, 1 \leq j \leq n}.$$

Since every  $(i, j) \in \{1, 2, \dots, n\}^2$  satisfies

$$\begin{aligned} & \sum_{k=1}^n \left( \delta_{i,k} - (1 - \delta_{i,n}) \delta_{f(i),k} \right) \frac{a_{k,j}}{a_{k,n}} \\ &= \underbrace{\sum_{k=1}^n \delta_{i,k} \frac{a_{k,j}}{a_{k,n}}}_{= \frac{a_{i,j}}{a_{i,n}}} - \sum_{k=1}^n \underbrace{(1 - \delta_{i,n}) \delta_{f(i),k} \frac{a_{k,j}}{a_{k,n}}}_{= (1 - \delta_{i,n}) \frac{a_{f(i),j}}{a_{f(i),n}}} \\ & \quad \text{(because the factor } \delta_{i,k} \text{ in the sum kills every addend except the one for } k=i) \quad \text{(because the factor } \delta_{f(i),k} \text{ in the sum kills every addend except the one for } k=f(i)) \\ &= \frac{a_{i,j}}{a_{i,n}} - \underbrace{(1 - \delta_{i,n})}_{=0} \frac{a_{f(i),j}}{a_{f(i),n}} = \frac{a_{i,j}}{a_{i,n}} - \begin{cases} 1, & \text{if } i < n; \frac{a_{f(i),j}}{a_{f(i),n}} \\ 0, & \text{if } i = n \end{cases} \\ &= \begin{cases} \frac{a_{i,j}}{a_{i,n}} - \frac{a_{f(i),j}}{a_{f(i),n}}, & \text{if } i < n; \\ \frac{a_{i,j}}{a_{i,n}}, & \text{if } i = n \end{cases} = d_{i,j} \quad \text{(by the definition of } d_{i,j}), \end{aligned}$$

this rewrites as

$$Z_f E = (d_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}.$$

Comparing this with  $D = (d_{i,j})_{1 \leq i \leq n, 1 \leq j \leq n}$  we obtain  $D = Z_f E$ . This proves (23).

Now, we can prove parts **(a)** and **(b)** of Theorem 2.9:

**(a)** Assume that the map  $f$  is not  $n$ -potent. Taking determinants on both sides of (23), we obtain

$$\det D = \det (Z_f E) = \underbrace{\det (Z_f)}_{=0} \cdot \det E = 0. \\ \text{(by Proposition 3.13 (b))}$$

Thus, (21) becomes

$$\det B = \left( \prod_{i=1}^{n-1} (a_{i,n} a_{f(i),n}) \right) \underbrace{\det D}_{=0} = 0.$$

This proves Theorem 2.9 **(a)**.

**(b)** Assume that the map  $f$  is  $n$ -potent. Taking determinants on both sides of



(23), we obtain

$$\begin{aligned}
 \det D = \det (Z_f E) &= \underbrace{\det (Z_f)}_{=1} \cdot \det E = \det E \\
 &\quad \text{(by Proposition 3.13 (a))} \\
 &= \underbrace{\left( \prod_{i=1}^n \frac{1}{a_{i,n}} \right)}_{\det A} \det A \quad \text{(by (22))} \\
 &= \left( \prod_{i=1}^{n-1} \frac{1}{a_{i,n}} \right) \cdot \frac{1}{a_{n,n}} \\
 &= \left( \prod_{i=1}^{n-1} \frac{1}{a_{i,n}} \right) \cdot \frac{1}{a_{n,n}} \det A.
 \end{aligned}$$

Thus, (21) becomes

$$\begin{aligned}
 \det B &= \left( \prod_{i=1}^{n-1} (a_{i,n} a_{f(i),n}) \right) \underbrace{\det D}_{= \left( \prod_{i=1}^{n-1} \frac{1}{a_{i,n}} \right) \cdot \frac{1}{a_{n,n}} \det A} \\
 &= \underbrace{\left( \prod_{i=1}^{n-1} (a_{i,n} a_{f(i),n}) \right)}_{= \prod_{i=1}^{n-1} a_{f(i),n} = \prod_{i \in \{1,2,\dots,n-1\}} a_{f(i),n}} \left( \prod_{i=1}^{n-1} \frac{1}{a_{i,n}} \right) \cdot \frac{1}{a_{n,n}} \det A \\
 &= \underbrace{\left( \prod_{i \in \{1,2,\dots,n-1\}} a_{f(i),n} \right)}_{= \frac{1}{a_{n,n}} \prod_{i \in \{1,2,\dots,n-1\}} a_{f(i),n} = \text{abut}_f A} \cdot \frac{1}{a_{n,n}} \det A = (\text{abut}_f A) \det A. \\
 &\quad \text{(by Remark 2.8 (a))}
 \end{aligned}$$

This proves Theorem 2.9 (b). □

### 3.6. Further questions

The above – rather indirect – road to the matrix-tree theorem suggests the following two questions:

- Is there a combinatorial proof of Theorem 2.9? Or, at least, is there a “division-free” proof (i.e., a proof that does not use a WLOG assumption that some of the  $a_{i,j}$  are invertible or a similar trick)?
- Can we similarly obtain some of the various generalizations and variants of the matrix-tree theorem, such as the all-minors matrix-tree theorem ([Chaiken82, (2)] and [Sahi13, Theorem 6])?

## 4. Appendix: some standard proofs

For the sake of completeness, let us give some proofs of standard results that have been used without proof above.

*Proof of Remark 2.6. (a)* We have  $1 \neq n$  (since  $n \geq 2$ ). But the map  $f$  is  $n$ -potent. Thus, there exists some  $k \in \mathbb{N}$  such that  $f^k(1) = n$ . Let  $h$  be the smallest such  $k$ . Then,  $f^h(1) = n$ . Hence,  $h \neq 0$  (since  $f^h(1) = n \neq 1 = f^0(1)$ ). Therefore,  $h - 1 \in \mathbb{N}$ , so that  $f^{h-1}(1) \neq n$  (because  $h$  is the **smallest**  $k \in \mathbb{N}$  such that  $f^k(1) = n$ ). Hence,  $f^{h-1}(1) \in \{1, 2, \dots, n-1\}$ . Thus,  $f^{h-1}(1)$  is a  $g \in \{1, 2, \dots, n-1\}$  such that  $f(g) = n$  (since  $f(f^{h-1}(1)) = f^h(1) = n$ ). Therefore, such a  $g$  exists. This proves Remark 2.6 (a).

**(b)** The map  $f$  is  $n$ -potent; thus,  $f(n) = n$ . Hence,  $n \in f^{-1}(n)$ . Remark 2.6 (a) shows that there exists some  $g \in \{1, 2, \dots, n-1\}$  such that  $f(g) = n$ . Consider this  $g$ . From  $f(g) = n$ , we obtain  $g \in f^{-1}(n)$ . From  $g \in \{1, 2, \dots, n-1\}$ , we obtain  $g \neq n$ . Hence,  $g$  and  $n$  are two distinct elements of the set  $f^{-1}(n)$ . Consequently,  $|f^{-1}(n)| \geq 2$ . This proves Remark 2.6 (b).  $\square$

*Proof of Remark 2.8. (b)* We have  $n \in f^{-1}(n)$  (since  $f(n) = n$ ) and  $g \in f^{-1}(n)$  (since  $f(g) = n$ ). Moreover,  $g \neq n$  (since  $g \in \{1, 2, \dots, n-1\}$ ). Hence,  $g$  and  $n$  are two distinct elements of  $f^{-1}(n)$ . Hence,  $|f^{-1}(n) \setminus \{n, g\}| = |f^{-1}(n)| - 2$ . But

$$\begin{aligned} & \{i \in \{1, 2, \dots, n-1\} \setminus \{g\} \mid f(i) = n\} \\ &= f^{-1}(n) \cap (\{1, 2, \dots, n-1\} \setminus \{g\}) \\ &= \underbrace{f^{-1}(n) \cap \{1, 2, \dots, n-1\}}_{=f^{-1}(n) \setminus \{n\}} \setminus \{g\} = (f^{-1}(n) \setminus \{n\}) \setminus \{g\} \\ &= f^{-1}(n) \setminus \{n, g\} \end{aligned}$$

so that

$$|\{i \in \{1, 2, \dots, n-1\} \setminus \{g\} \mid f(i) = n\}| = |f^{-1}(n) \setminus \{n, g\}| = |f^{-1}(n)| - 2. \quad (24)$$

Now,

$$\begin{aligned}
 & \prod_{\substack{i \in \{1, 2, \dots, n-1\}; \\ i \neq g}} a_{f(i), n} \\
 &= \prod_{i \in \{1, 2, \dots, n-1\} \setminus \{g\}} a_{f(i), n} = \left( \prod_{\substack{i \in \{1, 2, \dots, n-1\} \setminus \{g\}; \\ f(i) = n}} \underbrace{a_{f(i), n}}_{= a_{n, n}} \right) \left( \prod_{\substack{i \in \{1, 2, \dots, n-1\} \setminus \{g\}; \\ f(i) \neq n}} a_{f(i), n} \right) \\
 &= \left( \prod_{\substack{i \in \{1, 2, \dots, n-1\} \setminus \{g\}; \\ f(i) = n}} a_{n, n} \right) \left( \prod_{\substack{i \in \{1, 2, \dots, n-1\} \setminus \{g\}; \\ f(i) \neq n}} a_{f(i), n} \right) \\
 &= a_{n, n}^{|\{i \in \{1, 2, \dots, n-1\} \setminus \{g\} \mid f(i) = n\}|} a_{n, n}^{|\{f^{-1}(n)\}|-2} \\
 &\quad \text{(by (24))} \\
 &= a_{n, n}^{|\{f^{-1}(n)\}|-2} \left( \prod_{\substack{i \in \{1, 2, \dots, n-1\}; \\ f(i) \neq n}} a_{f(i), n} \right) = \text{abut}_f A
 \end{aligned}$$

(by the definition of  $\text{abut}_f A$ ). This proves Remark 2.8 (b).

(a) Assume that  $a_{n, n} \in \mathbb{K}$  is invertible. Fix  $g \in \{1, 2, \dots, n-1\}$  as in Remark 2.8 (b). Then,

$$\prod_{i \in \{1, 2, \dots, n-1\}} a_{f(i), n} = \underbrace{a_{f(g), n}}_{= a_{n, n}} \underbrace{\prod_{\substack{i \in \{1, 2, \dots, n-1\}; \\ i \neq g}} a_{f(i), n}}_{= \text{abut}_f A} = a_{n, n} \text{abut}_f A,$$

(by Remark 2.8 (b))

so that  $\text{abut}_f A = \frac{1}{a_{n, n}} \prod_{i \in \{1, 2, \dots, n-1\}} a_{f(i), n}$ . This proves Remark 2.8 (a). □

*Proof of Lemma 3.1.* We have  $G = \left( \sum_{k=1}^n b_{i, k} d_{i, j, k} \right)_{1 \leq i \leq m, 1 \leq j \leq m}$ . Thus, the definition of

a determinant yields

$$\begin{aligned}
 \det G &= \sum_{\sigma \in S_m} (-1)^\sigma \underbrace{\prod_{i=1}^m \left( \sum_{k=1}^n b_{i,k} d_{i,\sigma(i),k} \right)} \\
 &= \sum_{f: \{1,2,\dots,m\} \rightarrow \{1,2,\dots,n\}} \prod_{i=1}^m (b_{i,f(i)} d_{i,\sigma(i),f(i)}) \\
 &\quad \text{(by the product rule)} \\
 &= \sum_{\sigma \in S_m} (-1)^\sigma \sum_{f: \{1,2,\dots,m\} \rightarrow \{1,2,\dots,n\}} \prod_{i=1}^m (b_{i,f(i)} d_{i,\sigma(i),f(i)}) \\
 &= \sum_{f: \{1,2,\dots,m\} \rightarrow \{1,2,\dots,n\}} \sum_{\sigma \in S_m} (-1)^\sigma \underbrace{\prod_{i=1}^m (b_{i,f(i)} d_{i,\sigma(i),f(i)})}_{= \left( \prod_{i=1}^m b_{i,f(i)} \right) \left( \prod_{i=1}^m d_{i,\sigma(i),f(i)} \right)} \\
 &= \sum_{f: \{1,2,\dots,m\} \rightarrow \{1,2,\dots,n\}} \left( \prod_{i=1}^m b_{i,f(i)} \right) \underbrace{\sum_{\sigma \in S_m} (-1)^\sigma \left( \prod_{i=1}^m d_{i,\sigma(i),f(i)} \right)}_{= \det \left( (d_{i,j,f(i)})_{1 \leq i \leq m, 1 \leq j \leq m} \right)} \\
 &\quad \text{(by the definition of a determinant)} \\
 &= \sum_{f: \{1,2,\dots,m\} \rightarrow \{1,2,\dots,n\}} \left( \prod_{i=1}^m b_{i,f(i)} \right) \det \left( (d_{i,j,f(i)})_{1 \leq i \leq m, 1 \leq j \leq m} \right).
 \end{aligned}$$

□

*Proof of Proposition 3.3.* The elements  $f^0(i), f^1(i), \dots, f^n(i)$  are  $n + 1$  elements of the  $n$ -element set  $\{1, 2, \dots, n\}$ . Thus, by the pigeonhole principle, we see that two of these elements must be equal. In other words, there exist two elements  $u$  and  $v$  of  $\{0, 1, \dots, n\}$  such that  $u < v$  and  $f^u(i) = f^v(i)$ . Consider these  $u$  and  $v$ . We have  $v \in \{0, 1, \dots, n\}$ , so that  $v \leq n$  and thus  $v - 1 \leq n - 1$ . Hence,  $\{0, 1, \dots, v - 1\} \subseteq \{0, 1, \dots, n - 1\}$ .

We have  $u < v$ , so that  $u \leq v - 1$  (since  $u$  and  $v$  are integers). Thus,  $u \in \{0, 1, \dots, v - 1\}$  (since  $u$  is a nonnegative integer). Hence,  $0 \leq u \leq v - 1$ , so that  $0 \in \{0, 1, \dots, v - 1\}$ .

Let  $S$  be the set  $\{f^0(i), f^1(i), \dots, f^{v-1}(i)\}$ . From  $u \in \{0, 1, \dots, v - 1\}$ , we obtain  $f^u(i) \in \{f^0(i), f^1(i), \dots, f^{v-1}(i)\} = S$ . From  $0 \in \{0, 1, \dots, v - 1\}$ , we obtain  $f^0(i) \in \{f^0(i), f^1(i), \dots, f^{v-1}(i)\} = S$ .

Now,

$$f(s) \in S \quad \text{for every } s \in S \tag{25}$$

12.

---

<sup>12</sup>*Proof of (25):* Let  $s \in S$ .

We have  $s \in S = \{f^0(i), f^1(i), \dots, f^{v-1}(i)\}$ . In other words,  $s = f^h(i)$  for some  $h \in$

Now, we can easily see that

$$f^k(i) \in S \quad \text{for every } k \in \mathbb{N} \quad (26)$$

<sup>13</sup>.

On the other hand,

$$\begin{aligned} S &= \{f^0(i), f^1(i), \dots, f^{v-1}(i)\} = \{f^s(i) \mid s \in \{0, 1, \dots, v-1\}\} \\ &\subseteq \{f^s(i) \mid s \in \{0, 1, \dots, n-1\}\} \quad (\text{since } \{0, 1, \dots, v-1\} \subseteq \{0, 1, \dots, n-1\}). \end{aligned}$$

Hence, for every  $k \in \mathbb{N}$ , we have

$$\begin{aligned} f^k(i) &\in S \quad (\text{by (26)}) \\ &\subseteq \{f^s(i) \mid s \in \{0, 1, \dots, n-1\}\}. \end{aligned}$$

This proves Proposition 3.3. □

*Proof of Proposition 3.4.*  $\implies$ : Assume that  $f^{n-1}(i) = n$ . Thus, there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (namely,  $k = n - 1$ ). This proves the  $\implies$  direction of Proposition 3.4.

$\impliedby$ : Assume that there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$ . Consider this  $k$ . We must show that  $f^{n-1}(i) = n$ .

We have  $n = f^k(i) \in \{f^s(i) \mid s \in \{0, 1, \dots, n-1\}\}$  (by Proposition 3.3). In other words,  $n = f^s(i)$  for some  $s \in \{0, 1, \dots, n-1\}$ . Consider this  $s$ .

$\{0, 1, \dots, v-1\}$ . Consider this  $h$ . Thus,  $f \left( \underbrace{s}_{=f^h(i)} \right) = f(f^h(i)) = f^{h+1}(i)$ .

We want to prove that  $f(s) \in S$ . We are in one of the following two cases:

Case 1: We have  $h = v - 1$ .

Case 2: We have  $h \neq v - 1$ .

Let us first consider Case 1. In this case, we have  $h = v - 1$ . Hence,  $h + 1 = v$ . Now,  $f(s) = f^{h+1}(i) = f^v(i)$  (since  $h + 1 = v$ ). Compared with  $f^u(i) = f^v(i)$ , this yields  $f(s) = f^u(i) \in S$ . Hence,  $f(s) \in S$  is proven in Case 1.

Let us now consider Case 2. In this case, we have  $h \neq v - 1$ . Combined with  $h \in \{0, 1, \dots, v-1\}$ , this yields  $h \in \{0, 1, \dots, v-1\} \setminus \{v-1\} = \{0, 1, \dots, (v-1)-1\}$ , so that  $h + 1 \in \{0, 1, \dots, v-1\}$ . Thus,  $f^{h+1}(i) \in \{f^0(i), f^1(i), \dots, f^{v-1}(i)\} = S$ . Hence,  $f(s) = f^{h+1}(i) \in S$ . Thus,  $f(s) \in S$  is proven in Case 2.

We have now proven  $f(s) \in S$  in each of the two Cases 1 and 2. Thus,  $f(s) \in S$  always holds.

This proves (25).

<sup>13</sup>*Proof of (26):* We shall prove (26) by induction over  $k$ :

*Induction base:* We have  $f^0(i) \in S$ . In other words, (26) holds for  $k = 0$ . This completes the induction base.

*Induction step:* Let  $K \in \mathbb{N}$ . Assume that (26) holds for  $k = K$ . We must prove that (26) holds for  $k = K + 1$ .

We have assumed that (26) holds for  $k = K$ . In other words,  $f^K(i) \in S$ . Thus, (25) (applied to  $s = f^K(i)$ ) yields  $f(f^K(i)) \in S$ . Thus,  $f^{K+1}(i) = f(f^K(i)) \in S$ . In other words, (26) holds for  $k = K + 1$ . This completes the induction step. Hence, (26) is proven by induction.

We have  $f^s(i) = n$ . Using this fact (and the fact that  $f(n) = n$ ), we can prove (by induction over  $h$ ) that

$$f^h(i) = n \quad \text{for every integer } h \geq s. \quad (27)$$

But  $s \in \{0, 1, \dots, n-1\}$ , so that  $s \leq n-1$  and therefore  $n-1 \geq s$ . Hence, (27) (applied to  $h = n-1$ ) yields  $f^{n-1}(i) = n$ . This proves the  $\Leftarrow$  direction of Proposition 3.4.  $\square$

*Proof of Proposition 3.5.*  $\Leftarrow$ : Assume that  $f^{n-1}(\{1, 2, \dots, n\}) = \{n\}$ . For every  $i \in \{1, 2, \dots, n\}$ , we have

$$f^{n-1} \left( \underbrace{i}_{\in \{1, 2, \dots, n\}} \right) \in f^{n-1}(\{1, 2, \dots, n\}) = \{n\}$$

and thus  $f^{n-1}(i) = n$ . Hence, for every  $i \in \{1, 2, \dots, n-1\}$ , there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (namely,  $k = n-1$ ). In other words, the map  $f$  is  $n$ -potent. This proves the  $\Leftarrow$  direction of Proposition 3.5.

$\Rightarrow$ : Assume that the map  $f$  is  $n$ -potent. Let  $i \in \{1, 2, \dots, n-1\}$ . Then, there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (since  $f$  is  $n$ -potent). Thus,  $f^{n-1}(i) = n$  (by the  $\Leftarrow$  direction of Proposition 3.4).

Now, forget that we fixed  $i$ . We thus have shown that  $f^{n-1}(i) = n$  for each  $i \in \{1, 2, \dots, n\}$ . Hence,

$$\{f^{n-1}(1), f^{n-1}(2), \dots, f^{n-1}(n)\} = \left\{ \underbrace{n, n, \dots, n}_{n \text{ times } n} \right\} = \{n\}.$$

Thus,  $f^{n-1}(\{1, 2, \dots, n\}) = \{f^{n-1}(1), f^{n-1}(2), \dots, f^{n-1}(n)\} = \{n\}$ . This proves the  $\Rightarrow$  direction of Proposition 3.5.  $\square$

*Proof of Corollary 3.6.* We are in one of the following two cases:

Case 1: We have  $f^{n-1}(i) = n$ .

Case 2: We have  $f^{n-1}(i) \neq n$ .

Let us consider Case 1 first. In this case, we have  $f^{n-1}(i) = n$ . Thus,  $\delta_{f^{n-1}(i), n} = 1$ .

But  $f^n(i) = f \left( \underbrace{f^{n-1}(i)}_{=n} \right) = f(n) = n$ , so that  $\delta_{f^n(i), n} = 1$ . Hence,  $\delta_{f^{n-1}(i), n} = 1 = \delta_{f^n(i), n}$ . Thus, Corollary 3.6 is proven in Case 1.

Let us now consider Case 2. In this case, we have  $f^{n-1}(i) \neq n$ . Thus,  $\delta_{f^{n-1}(i), n} = 0$ . On the other hand, we have  $f^n(i) \neq n$ <sup>14</sup>. Hence,  $\delta_{f^n(i), n} = 0$ . Hence,  $\delta_{f^{n-1}(i), n} = 0 = \delta_{f^n(i), n}$ . Thus, Corollary 3.6 is proven in Case 2.

---

<sup>14</sup>*Proof.* Assume the contrary. Thus,  $f^n(i) = n$ . Hence, there exists some  $k \in \mathbb{N}$  such that  $f^k(i) = n$  (namely,  $k = n$ ). Thus,  $f^{n-1}(i) = n$  (according to the  $\Leftarrow$  direction of Proposition 3.4). This contradicts  $f^{n-1}(i) \neq n$ . This contradiction proves that our assumption was wrong, qed.

Now, we have proven Corollary 3.6 in each of the two Cases 1 and 2. Hence, Corollary 3.6 always holds.  $\square$

*Proof of Lemma 3.14.* The definition of  $\det A$  yields  $\det A = \sum_{\sigma \in S_m} (-1)^\sigma \prod_{i=1}^m a_{i,\sigma(i)}$  (since  $A = (a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq m}$ ). On the other hand, the definition of  $\det \left( (b_i a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq m} \right)$  yields

$$\begin{aligned} \det \left( (b_i a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq m} \right) &= \sum_{\sigma \in S_m} (-1)^\sigma \underbrace{\prod_{i=1}^m (b_i a_{i,\sigma(i)})}_{= \left( \prod_{i=1}^m b_i \right) \left( \prod_{i=1}^m a_{i,\sigma(i)} \right)} \\ &= \sum_{\sigma \in S_m} (-1)^\sigma \left( \prod_{i=1}^m b_i \right) \left( \prod_{i=1}^m a_{i,\sigma(i)} \right) \\ &= \left( \prod_{i=1}^m b_i \right) \underbrace{\sum_{\sigma \in S_m} (-1)^\sigma \prod_{i=1}^m a_{i,\sigma(i)}}_{=\det A} = \left( \prod_{i=1}^m b_i \right) \det A. \end{aligned}$$

This proves Lemma 3.14.  $\square$

## References

- [Abeles14] Francine F. Abeles, *Chio's and Dodgson's determinantal identities*, Linear Algebra and its Applications, Volume 454, 1 August 2014, pp. 130–137.
- [BerBru08] Adam Berliner and Richard A. Brualdi, *A combinatorial proof of the Dodgson/Muir determinantal identity*, International Journal of Information and Systems Sciences, Volume 4 (2008), Number 1, pp. 1–7.
- [Chaiken82] Seth Chaiken, *A combinatorial proof of the all minors matrix tree theorem*, SIAM J. Alg. Disc. Math., Vol. 3, No. 3, September 1982, pp. 319–329.
- [Conrad09] Keith Conrad, *Universal identities*, 12 October 2009.  
<http://www.math.uconn.edu/~kconrad/blurbs/linmultialg/univid.pdf>
- [Eves68] Howard Eves, *Elementary Matrix Theory*, Allyn & Bacon, 2nd printing 1968.
- [Grinbe15] Darij Grinberg, *Notes on the combinatorial fundamentals of algebra*, 10 January 2019.  
<http://www.cip.ifi.lmu.de/~grinberg/primes2015/sols.pdf>  
The numbering of theorems and formulas in this link might shift when

the project gets updated; for a “frozen” version whose numbering matches that in the citations above, see <https://github.com/darijgr/detnotes/releases/tag/2019-01-10>.

- [Heinig11] Peter Christian Heinig, *Chio Condensation and Random Sign Matrices*, arXiv:1103.2717v3.
  - [KarZha16] Karthik Karnik, Anya Zhang, *Combinatorial proof of Chio Pivotal Condensation*, 25 May 2016.
  - [Sahi13] Siddhartha Sahi, *Harmonic vectors and matrix tree theorems*, arXiv:1309.4047v1.
  - [Verstr12] Jacques Verstraete, *Math264A Lecture J*, 4 December 2012.  
<http://www.math.ucsd.edu/~jverstra/264A-LECTUREJ.pdf>
  - [Zeilbe85] Doron Zeilberger, *A combinatorial approach to matrix algebra*, *Discrete Mathematics* 56 (1985), pp. 61–72.
-