

Noncommutative birational rowmotion on a rectangle

A case study in noncommutative dynamics

Darij Grinberg (Drexel University)
joint work with Tom Roby (UConn)

29 March 2023, Kungliga Tekniska Högskolan, Stockholm

slides: [http:](http://www.cip.ifi.lmu.de/~grinberg/algebra/kth2023.pdf)

[//www.cip.ifi.lmu.de/~grinberg/algebra/kth2023.pdf](http://www.cip.ifi.lmu.de/~grinberg/algebra/kth2023.pdf)


paper: <https://arxiv.org/abs/2208.11156>

FPSAC abstract: [https:](https://www.cip.ifi.lmu.de/~grinberg/algebra/fps2023.pdf)

[//www.cip.ifi.lmu.de/~grinberg/algebra/fps2023.pdf](https://www.cip.ifi.lmu.de/~grinberg/algebra/fps2023.pdf)

Introduction: Posets

- A **poset** (= partially ordered set) is a set P with a reflexive, transitive and antisymmetric relation.
- We use the symbols $<$, \leq , $>$ and \geq accordingly.
- We draw posets as Hasse diagrams:




- We only care about finite posets here.
- We say that $u \in P$ is **covered by** $v \in P$ (written $u \triangleleft v$) if we have $u < v$ and there is no $w \in P$ satisfying $u < w < v$.
- We say that $u \in P$ **covers** $v \in P$ (written $u \triangleright v$) if we have $u > v$ and there is no $w \in P$ satisfying $u > w > v$.


More poset basics: \widehat{P}

- Let P be a finite poset. We define \widehat{P} to be the poset obtained by adjoining two new elements 0 and 1 to P and forcing
 - 0 to be less than every other element, and
 - 1 to be greater than every other element.

Example:



- A **linear extension** of P means a list (v_1, v_2, \dots, v_n) of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- For instance,





has two linear extensions $(\alpha, \beta, \gamma, \delta)$ and $(\beta, \alpha, \gamma, \delta)$.

- Every finite poset has at least one linear extension.

More poset basics: order ideals

- An **order ideal** of a poset P is a subset S of P such that if $v \in S$ and $w \leq v$, then $w \in S$.
- Examples (the elements of the order ideal are marked in red):



- We let $J(P)$ denote the set of all order ideals of P .

- **Classical rowmotion** is the rowmotion studied by Striker/Williams ([arXiv:1108.1172](https://arxiv.org/abs/1108.1172)). It has appeared many times before, under different guises:
 - Brouwer/Schrijver (1974) (as a permutation of the antichains),
 - Fon-der-Flaass (1993) (as a permutation of the antichains),
 - Cameron/Fon-der-Flaass (1995) (as a permutation of the monotone Boolean functions),
 - Panyushev (2008), Armstrong-Stump-Thomas (2011) (as a permutation of the antichains or “nonnesting partitions”, with relations to Lie theory).


Classical rowmotion: the standard definition

- Let P be a finite poset. **Classical rowmotion** is the map $\mathbf{r} : J(P) \rightarrow J(P)$ which sends every order ideal S to a new order ideal $\mathbf{r}(S)$ defined as follows:
 - Invert colors** (i.e., take the complement $P \setminus S$).
 - Boil down to generators** (i.e., take the set M of minimal elements of this complement).
 - Complete downwards** (i.e., take the set J of all $w \in P$ such that there exists an $m \in M$ such that $w \leq m$).

Then, $\mathbf{r}(S) = J$.

Example:

Let S be the following order ideal (\bullet = inside order ideal):




Classical rowmotion: the standard definition

- Let P be a finite poset. **Classical rowmotion** is the map $\mathbf{r} : J(P) \rightarrow J(P)$ which sends every order ideal S to a new order ideal $\mathbf{r}(S)$ defined as follows:
 - **Invert colors** (i.e., take the complement $P \setminus S$).
 - **Boil down to generators** (i.e., take the set M of minimal elements of this complement).
 - **Complete downwards** (i.e., take the set J of all $w \in P$ such that there exists an $m \in M$ such that $w \leq m$).

Then, $\mathbf{r}(S) = J$.

Example:

Mark the elements of the complement blue.




Classical rowmotion: the standard definition

- Let P be a finite poset. **Classical rowmotion** is the map $\mathbf{r} : J(P) \rightarrow J(P)$ which sends every order ideal S to a new order ideal $\mathbf{r}(S)$ defined as follows:
 - **Invert colors** (i.e., take the complement $P \setminus S$).
 - **Boil down to generators** (i.e., take the set M of minimal elements of this complement).
 - **Complete downwards** (i.e., take the set J of all $w \in P$ such that there exists an $m \in M$ such that $w \leq m$).

Then, $\mathbf{r}(S) = J$.

Example:

Leave only the minimal elements:




Classical rowmotion: the standard definition

- Let P be a finite poset. **Classical rowmotion** is the map $\mathbf{r} : J(P) \rightarrow J(P)$ which sends every order ideal S to a new order ideal $\mathbf{r}(S)$ defined as follows:
 - **Invert colors** (i.e., take the complement $P \setminus S$).
 - **Boil down to generators** (i.e., take the set M of minimal elements of this complement).
 - **Complete downwards** (i.e., take the set J of all $w \in P$ such that there exists an $m \in M$ such that $w \leq m$).

Then, $\mathbf{r}(S) = J$.

Example:

$\mathbf{r}(S)$ is the order ideal generated by M (“everything below M ”):



Classical rowmotion: properties

Classical rowmotion is a permutation of $J(P)$, hence has finite order. This order can be fairly large.


Classical rowmotion: properties

Classical rowmotion is a permutation of $J(P)$, hence has finite order. This order can be fairly large.

However, **for some types of P** , the order can be explicitly computed or bounded from above.

See Striker/Williams ([arXiv:1108.1172](#)) for the first generation of results.


- If P is a $p \times q$ -rectangle:



(shown here for $p = 2$ and $q = 3$), then $\text{ord}(\mathbf{r}) = p + q$.


Example:

Let S be the order ideal of the 2×3 -rectangle given by:




Example:

$r(S)$ is




Example:

$r^2(S)$ is




Example:

$r^3(S)$ is




Example:

$r^4(S)$ is



Example:

$r^5(S)$ is




which is precisely the S we started with.

$$\text{ord}(\mathbf{r}) = p + q = 2 + 3 = 5.$$

Further posets for which classical rowmotion has small order:

- If P is a Δ -shaped triangle with sidelength $p - 1$:




(shown here for $p = 4$), then $\text{ord}(\mathbf{r}) = 2p$ (if $p > 2$).

- In this case, \mathbf{r}^p is “reflection in the y -axis” (i.e., the central vertical axis).

Further posets for which classical rowmotion has small order:

- If P is a Δ -shaped triangle with sidelength $p - 1$:



(shown here for $p = 4$), then $\text{ord}(\mathbf{r}) = 2p$ (if $p > 2$).

- In this case, \mathbf{r}^P is “reflection in the y -axis” (i.e., the central vertical axis).
- More general examples come from finite Weyl groups (Armstrong/Stump/Thomas, [arXiv:1101.1277](#)) and from minuscule weights of classical groups (Rush/Shi, [arXiv:1108.5245](#); Okada, [arXiv:2004.05364](#)).

There is an alternative definition of classical rowmotion, which splits it into many little steps.

- If P is a poset and $v \in P$, then the v -**toggle** is the map $\mathbf{t}_v : J(P) \rightarrow J(P)$ which takes every order ideal S to:
 - $S \cup \{v\}$, if v is not in S but all elements of P covered by v are in S already;
 - $S \setminus \{v\}$, if v is in S but none of the elements of P covering v is in S ;
 - S otherwise.
- Simpler way to state this: $\mathbf{t}_v(S)$ is:
 - $S \Delta \{v\}$ (symmetric difference) if this is an order ideal;
 - S otherwise.

(“Try to add or remove v from S ; if this breaks the order ideal axiom, leave S fixed.”)


Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

Start with this order ideal S :




Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

First apply $\mathbf{t}_{(2,2)}$, which changes nothing:




Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

Then apply $\mathbf{t}_{(1,2)}$, which adds $(1,2)$ to the order ideal:




Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

Then apply $\mathbf{t}_{(2,1)}$, which removes $(2, 1)$ from the order ideal:




Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

Finally apply $\mathbf{t}_{(1,1)}$, which changes nothing:




Classical rowmotion: the toggling definition

- Let (v_1, v_2, \dots, v_n) be a **linear extension** of P ; this means a list of all elements of P (each only once) such that $i < j$ whenever $v_i < v_j$.
- Cameron and Fon-der-Flaass showed that

$$\mathbf{r} = \mathbf{t}_{v_1} \circ \mathbf{t}_{v_2} \circ \dots \circ \mathbf{t}_{v_n}.$$

Example:

So this is $\mathbf{r}(S)$:




- define **noncommutative birational rowmotion**: a generalization of classical rowmotion on several levels, due to David Einstein, James Propp, Tom Roby and myself, based on ideas of Anatol Kirillov and Arkady Berenstein.
- extend the “order $p + q$ ” theorem for rectangles to this generalization.
- ask some questions.

Noncommutative birational rowmotion: definition

- Let \mathbb{K} be a ring (not necessarily commutative).
- A \mathbb{K} -labelling of P will mean a function $\widehat{P} \rightarrow \mathbb{K}$.
- The values of such a function will be called the **labels** of the labelling.
- We will represent labellings by drawing the labels on the vertices of the Hasse diagram of \widehat{P} .

Example: This is a \mathbb{Q} -labelling of the 2×2 -rectangle:



- For any $v \in P$, define the **birational v -toggle** as the partial map $T_v : \mathbb{K}^{\widehat{P}} \dashrightarrow \mathbb{K}^{\widehat{P}}$ defined by

$$(T_v f)(w) = \begin{cases} f(w), & \text{if } w \neq v; \\ \left(\sum_{\substack{u \in \widehat{P}; \\ u < v}} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{\substack{u \in \widehat{P}; \\ u > v}} \overline{f(u)}}, & \text{if } w = v \end{cases}$$

for all $w \in \widehat{P}$.

Here (and in the following), \overline{m} means m^{-1} whenever $m \in \mathbb{K}$.

- For any $v \in P$, define the **birational v -toggle** as the partial map $T_v : \mathbb{K}^{\hat{P}} \dashrightarrow \mathbb{K}^{\hat{P}}$ defined by

$$(T_v f)(w) = \begin{cases} f(w), & \text{if } w \neq v; \\ \left(\sum_{\substack{u \in \hat{P}; \\ u < v}} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{\substack{u \in \hat{P}; \\ u > v}} \overline{f(u)}}, & \text{if } w = v \end{cases}$$

for all $w \in \hat{P}$.

Here (and in the following), \overline{m} means m^{-1} whenever $m \in \mathbb{K}$.

- This is a **partial** map. If any of the inverses does not exist in \mathbb{K} , then $T_v f$ is undefined!

- For any $v \in P$, define the **birational v -toggle** as the partial map $T_v : \mathbb{K}^{\hat{P}} \dashrightarrow \mathbb{K}^{\hat{P}}$ defined by

$$(T_v f)(w) = \begin{cases} f(w), & \text{if } w \neq v; \\ \left(\sum_{\substack{u \in \hat{P}; \\ u < v}} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{\substack{u \in \hat{P}; \\ u > v}} f(u)}, & \text{if } w = v \end{cases}$$

for all $w \in \hat{P}$.

Here (and in the following), \overline{m} means m^{-1} whenever $m \in \mathbb{K}$.

- This is a **partial** map. If any of the inverses does not exist in \mathbb{K} , then $T_v f$ is undefined!
- Notice that this is a **local change** to the label at v ; all other labels stay the same.

- For any $v \in P$, define the **birational v -toggle** as the partial map $T_v : \mathbb{K}^{\widehat{P}} \dashrightarrow \mathbb{K}^{\widehat{P}}$ defined by

$$(T_v f)(w) = \begin{cases} f(w), & \text{if } w \neq v; \\ \left(\sum_{\substack{u \in \widehat{P}; \\ u < v}} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{\substack{u \in \widehat{P}; \\ u > v}} f(u)}, & \text{if } w = v \end{cases}$$

for all $w \in \widehat{P}$.

Here (and in the following), \overline{m} means m^{-1} whenever $m \in \mathbb{K}$.

- This is a **partial** map. If any of the inverses does not exist in \mathbb{K} , then $T_v f$ is undefined!
- Notice that this is a **local change** to the label at v ; all other labels stay the same.
- If \mathbb{K} is commutative, then $T_v^2 = \text{id}$ (on the range of T_v).

- We define **(noncommutative) birational rowmotion** as the partial map

$$R := T_{v_1} \circ T_{v_2} \circ \cdots \circ T_{v_n} : \mathbb{K}^{\hat{P}} \dashrightarrow \mathbb{K}^{\hat{P}},$$

where (v_1, v_2, \dots, v_n) is a linear extension of P .

- This is indeed independent on the linear extension, because:

- We define **(noncommutative) birational rowmotion** as the partial map


$$R := T_{v_1} \circ T_{v_2} \circ \cdots \circ T_{v_n} : \mathbb{K}^{\hat{P}} \dashrightarrow \mathbb{K}^{\hat{P}},$$

where (v_1, v_2, \dots, v_n) is a linear extension of P .

- This is indeed independent on the linear extension, because:
 - T_v and T_w commute whenever v and w are incomparable (or just don't cover each other);
 - we can get from any linear extension to any other by switching incomparable adjacent elements.


Example:

Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:



Example:

Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:




We have $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$ (using the linear extension $((1, 1), (1, 2), (2, 1), (2, 2))$).

That is, toggle in the order “top, left, right, bottom”.

Example:


Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:



We are using $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$.

Example:


Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:



We are using $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$.

Example:



Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:



We are using $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$.

Example:


Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:

original labelling f	labelling $T_{(1,1)} T_{(1,2)} T_{(2,1)} T_{(2,2)} f = Rf$
	

We are using $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$.

Example:

Let us “rowmote” a (generic) \mathbb{K} -labelling of the 2×2 -rectangle:



We have used $R = T_{(1,1)} \circ T_{(1,2)} \circ T_{(2,1)} \circ T_{(2,2)}$ and simplified the result.

Birational rowmotion: motivation

- Why is this called birational rowmotion?
- Indeed, it generalizes classical rowmotion of order ideals:
 - Let $\text{Trop } \mathbb{Z}$ be the **tropical semiring** over \mathbb{Z} . This is the set $\mathbb{Z} \cup \{-\infty\}$ with “addition” $(a, b) \mapsto \max\{a, b\}$ and “multiplication” $(a, b) \mapsto a + b$. This is a semifield.

- Why is this called birational rowmotion?
- Indeed, it generalizes classical rowmotion of order ideals:
 - Let $\text{Trop } \mathbb{Z}$ be the **tropical semiring** over \mathbb{Z} . This is the set $\mathbb{Z} \cup \{-\infty\}$ with “addition” $(a, b) \mapsto \max\{a, b\}$ and “multiplication” $(a, b) \mapsto a + b$. This is a semifield.
 - To every order ideal $S \in J(P)$, assign a $\text{Trop } \mathbb{Z}$ -labelling $\text{tlab } S$ defined by

$$(\text{tlab } S)(v) = \begin{cases} 1, & \text{if } v \notin S \cup \{0\}; \\ 0, & \text{if } v \in S \cup \{0\}. \end{cases}$$

This map $\text{tlab} : J(P) \rightarrow (\text{Trop } \mathbb{Z})^{\hat{P}}$ is injective.

- Why is this called birational rowmotion?
- Indeed, it generalizes classical rowmotion of order ideals:
 - Let $\text{Trop } \mathbb{Z}$ be the **tropical semiring** over \mathbb{Z} . This is the set $\mathbb{Z} \cup \{-\infty\}$ with “addition” $(a, b) \mapsto \max\{a, b\}$ and “multiplication” $(a, b) \mapsto a + b$. This is a semifield.
 - To every order ideal $S \in J(P)$, assign a $\text{Trop } \mathbb{Z}$ -labelling $\text{tlab } S$ defined by

$$(\text{tlab } S)(v) = \begin{cases} 1, & \text{if } v \notin S \cup \{0\}; \\ 0, & \text{if } v \in S \cup \{0\}. \end{cases}$$

This map $\text{tlab} : J(P) \rightarrow (\text{Trop } \mathbb{Z})^{\hat{P}}$ is injective.

- Let \mathbf{t}_v be the order ideal v -toggle, and let \mathbf{r} be order ideal rowmotion. Then:

$$T_v \circ \text{tlab} = \text{tlab} \circ \mathbf{t}_v, \quad R \circ \text{tlab} = \text{tlab} \circ \mathbf{r}.$$

Birational rowmotion: motivation

- Why is this called birational rowmotion?
- Indeed, it generalizes classical rowmotion of order ideals:
 - Let $\text{Trop } \mathbb{Z}$ be the **tropical semiring** over \mathbb{Z} . This is the set $\mathbb{Z} \cup \{-\infty\}$ with “addition” $(a, b) \mapsto \max\{a, b\}$ and “multiplication” $(a, b) \mapsto a + b$. This is a semifield.
 - To every order ideal $S \in J(P)$, assign a $\text{Trop } \mathbb{Z}$ -labelling $\text{tlab } S$ defined by

$$(\text{tlab } S)(v) = \begin{cases} 1, & \text{if } v \notin S \cup \{0\}; \\ 0, & \text{if } v \in S \cup \{0\}. \end{cases}$$

This map $\text{tlab} : J(P) \rightarrow (\text{Trop } \mathbb{Z})^{\hat{P}}$ is injective.

- Let \mathbf{t}_v be the order ideal v -toggle, and let \mathbf{r} be order ideal rowmotion. Then:

$$T_v \circ \text{tlab} = \text{tlab} \circ \mathbf{t}_v, \quad R \circ \text{tlab} = \text{tlab} \circ \mathbf{r}.$$

- Don't like semifields? Use \mathbb{Q} and take the “tropical limit”.

- If \mathbb{K} is commutative, then birational rowmotion R has nice orders for nice posets (mostly [Grinberg/Roby 2014](#)):
 - If P is a rectangle $[p] \times [q]$, then $R^{p+q} = \text{id}$.

- If \mathbb{K} is commutative, then birational rowmotion R has nice orders for nice posets (mostly [Grinberg/Roby 2014](#)):
 - If P is a rectangle $[p] \times [q]$, then $R^{p+q} = \text{id}$.
 - If P is a “right half” \triangleright of the square $[p] \times [p]$, then $R^{2p} = \text{id}$.


- If \mathbb{K} is commutative, then birational rowmotion R has nice orders for nice posets (mostly [Grinberg/Roby 2014](#)):
 - If P is a rectangle $[p] \times [q]$, then $R^{p+q} = \text{id}$.
 - If P is a “right half” \triangleright of the square $[p] \times [p]$, then $R^{2p} = \text{id}$.
 - If P is a “top half” Δ or “bottom half” ∇ of the square $[p] \times [p]$, then $R^{2p} = \text{id}$, and moreover R^p is reflection across the vertical axis.

- If \mathbb{K} is commutative, then birational rowmotion R has nice orders for nice posets (mostly [Grinberg/Roby 2014](#)):
 - If P is a rectangle $[p] \times [q]$, then $R^{p+q} = \text{id}$.
 - If P is a “right half” \triangleright of the square $[p] \times [p]$, then $R^{2p} = \text{id}$.
 - If P is a “top half” Δ or “bottom half” ∇ of the square $[p] \times [p]$, then $R^{2p} = \text{id}$, and moreover R^p is reflection across the vertical axis.
 - More generally, if P is the minuscule poset associated to a minuscule weight λ of a finite-dimensional simple Lie algebra \mathfrak{g} , then $R^h = \text{id}$, where h is the Coxeter number of \mathfrak{g} . ([Soichi Okada, doi:10.37236/9557](#) .)

- If \mathbb{K} is commutative, then birational rowmotion R has nice orders for nice posets (mostly [Grinberg/Roby 2014](#)):
 - If P is a rectangle $[p] \times [q]$, then $R^{p+q} = \text{id}$.
 - If P is a “right half” \triangleright of the square $[p] \times [p]$, then $R^{2p} = \text{id}$.
 - If P is a “top half” Δ or “bottom half” ∇ of the square $[p] \times [p]$, then $R^{2p} = \text{id}$, and moreover R^p is reflection across the vertical axis.
 - More generally, if P is the minuscule poset associated to a minuscule weight λ of a finite-dimensional simple Lie algebra \mathfrak{g} , then $R^h = \text{id}$, where h is the Coxeter number of \mathfrak{g} . ([Soichi Okada, doi:10.37236/9557](#) .)
 - If P is an “ n -graded forest” (a forest with all leaves having rank n), then $R^\ell = \text{id}$ for $\ell = \text{lcm}(1, 2, \dots, n + 1)$.


Birational rowmotion: some chaos

- In general, even if \mathbb{K} is commutative, R can have infinite order
– e.g., for the following two posets:



Birational rowmotion: some chaos


- In general, even if \mathbb{K} is commutative, R can have infinite order
– e.g., for the following two posets:




- Things get even more complicated when \mathbb{K} is noncommutative...

Birational rowmotion: some chaos

- In general, even if \mathbb{K} is commutative, R can have infinite order
– e.g., for the following two posets:




- Things get even more complicated when \mathbb{K} is noncommutative...
- Take this poset:




This satisfies $R^6 = \text{id}$ if \mathbb{K} is commutative, but nothing like that in general.

Birational rowmotion: some chaos

- In general, even if \mathbb{K} is commutative, R can have infinite order
– e.g., for the following two posets:



- Things get even more complicated when \mathbb{K} is noncommutative...
- Take this poset:



This satisfies $R^6 = \text{id}$ if \mathbb{K} is commutative, but nothing like that in general.

- However, not all is lost!


Birational rowmotion: the rectangle case

- Let p and q be two positive integers. Let \mathbb{K} be a ring. Let P be the $p \times q$ -rectangle poset: i.e.,

$$P := [p] \times [q], \quad \text{where } [m] := \{1, 2, \dots, m\}.$$

(The order on P is entrywise.)

Example: For $p = 3$ and $q = 4$, this is



- Let p and q be two positive integers. Let \mathbb{K} be a ring. Let P be the $p \times q$ -rectangle poset: i.e.,

$$P := [p] \times [q], \quad \text{where } [m] := \{1, 2, \dots, m\}.$$

(The order on P is entrywise.)

- Let $f \in \mathbb{K}^{\hat{P}}$ be a \mathbb{K} -labelling. Let $a = f(0)$ and $b = f(1)$.

Birational rowmotion: the rectangle case

- Let p and q be two positive integers. Let \mathbb{K} be a ring. Let P be the $p \times q$ -rectangle poset: i.e.,

$$P := [p] \times [q], \quad \text{where } [m] := \{1, 2, \dots, m\}.$$

(The order on P is entrywise.)

- Let $f \in \mathbb{K}^{\hat{P}}$ be a \mathbb{K} -labelling. Let $a = f(0)$ and $b = f(1)$.

Periodicity theorem (* 2015, † 2022 G & Roby):

If a and b are invertible and $R^{p+q}f$ is well-defined, then

$$(R^{p+q}f)(x) = a\bar{b} \cdot f(x) \cdot \bar{a}b \quad \text{for each } x \in \hat{P}.$$

Note that $a\bar{b} \cdot f(x) \cdot \bar{a}b$ is **not** generally conjugate to $f(x)$.

Birational rowmotion: the rectangle case

- Let p and q be two positive integers. Let \mathbb{K} be a ring. Let P be the $p \times q$ -rectangle poset: i.e.,

$$P := [p] \times [q], \quad \text{where } [m] := \{1, 2, \dots, m\}.$$

(The order on P is entrywise.)

- Let $f \in \mathbb{K}^{\hat{P}}$ be a \mathbb{K} -labelling. Let $a = f(0)$ and $b = f(1)$.

Periodicity theorem (* 2015, † 2022 G & Roby):

If a and b are invertible and $R^{p+q}f$ is well-defined, then

$$(R^{p+q}f)(x) = a\bar{b} \cdot f(x) \cdot \bar{a}b \quad \text{for each } x \in \hat{P}.$$


Reciprocity theorem (* 2015, † 2022 G & Roby):

Let $\ell \in \mathbb{N}$. Let $(i, j) \in P$. If $R^\ell f$ is well-defined and $\ell \geq i + j - 1$, then

$$(R^\ell f)(i, j) = a \cdot \overbrace{(R^{\ell-i-j+1}f)(p+1-i, q+1-j)}^{\text{=antipode of } (i,j) \text{ in } P} \cdot b.$$


- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^0 f$:




- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^1 f$:




- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^2 f$:




- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^3 f$:




- Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^4 f$:



- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.


Here is $R^4 f$:



(after nontrivial simplifications).

- **Example:** Iteratively apply R to a labelling of the 2×2 -rectangle.

Here is $R^4 f$:



This confirms the periodicity theorem for $p = q = 2$.

- Note that this is similar to Kontsevich's periodicity conjecture, proved by Lyudu/Shkarin ([arXiv:1305.1965](https://arxiv.org/abs/1305.1965)).

Birational rowmotion: the rectangle case, example

- Here are $R^0 f, R^1 f, \dots, R^4 f$ for a generic $f \in \widehat{\mathbb{K}^{[2] \times [2]}}$ again, this time fully simplified and with the $f(0) = a$ and $f(1) = b$ labels removed:

$$R^0 f = \begin{array}{ccc} & z & \\ x & / \quad \backslash & y \\ & w & \end{array} ; \quad R^1 f = \begin{array}{ccc} & (x+y)\bar{z}b & \\ w\bar{x}(x+y)\bar{z}b & / \quad \backslash & w\bar{y}(x+y)\bar{z}b \\ & a\bar{z}b & \end{array}$$

$$R^2 f = \begin{array}{ccc} & w(\bar{x} + \bar{y})b & \\ a\bar{y}b & / \quad \backslash & a\bar{x}b \\ & a\bar{b}z\bar{x} + \bar{y}b & \end{array} ; \quad R^3 f = \begin{array}{ccc} & a\bar{w}b & \\ a\bar{b}z\bar{x} + \bar{y}w\bar{b} & / \quad \backslash & a\bar{b}z\bar{x} + \bar{y}x\bar{w}b \\ & a\bar{b} \cdot \bar{x} + \bar{y} \cdot \bar{w}b & \end{array}$$

Birational rowmotion: the rectangle case, example

- Here are $R^0 f, R^1 f, \dots, R^4 f$ for a generic $f \in \widehat{\mathbb{K}^{[2] \times [2]}}$ again, this time fully simplified and with the $f(0) = a$ and $f(1) = b$ labels removed:

$$R^0 f = \begin{array}{ccc} & z & \\ & / \quad \backslash & \\ x & & y \\ & \backslash \quad / & \\ & w & \end{array} ; \quad R^1 f = \begin{array}{ccc} & (x+y)\bar{z}b & \\ & / \quad \backslash & \\ w\bar{x}(x+y)\bar{z}b & & w\bar{y}(x+y)\bar{z}b \\ & \backslash \quad / & \\ & a\bar{z}b & \end{array}$$

$$R^2 f = \begin{array}{ccc} & w(\bar{x} + \bar{y})b & \\ & / \quad \backslash & \\ a\bar{y}b & & a\bar{x}b \\ & \backslash \quad / & \\ & a\bar{b}z\bar{x} + \bar{y}b & \end{array} ; \quad R^3 f = \begin{array}{ccc} & a\bar{w}b & \\ & / \quad \backslash & \\ a\bar{b}z\bar{x} + \bar{y}w\bar{b} & & a\bar{b}z\bar{x} + \bar{y}x\bar{w}b \\ & \backslash \quad / & \\ & a\bar{b} \cdot \bar{x} + \bar{y} \cdot \bar{w}b & \end{array}$$

Equally colored labels are related by reciprocity. Can you spot some more?

Birational rowmotion: the rectangle case, example

- Here are $R^0 f, R^1 f, \dots, R^4 f$ for a generic $f \in \widehat{\mathbb{K}^{[2] \times [2]}}$ again, this time fully simplified and with the $f(0) = a$ and $f(1) = b$ labels removed:

$$R^0 f = \begin{array}{ccc} & z & \\ x & / \quad \backslash & y \\ & w & \end{array} ; \quad R^1 f = \begin{array}{ccc} & (x+y)\bar{z}b & \\ w\bar{x}(x+y)\bar{z}b & / \quad \backslash & w\bar{y}(x+y)\bar{z}b \\ & a\bar{z}b & \end{array}$$

$$R^2 f = \begin{array}{ccc} & w(\bar{x} + \bar{y})b & \\ a\bar{y}b & / \quad \backslash & a\bar{x}b \\ & \overline{abzx + yb} & \end{array} ; \quad R^3 f = \begin{array}{ccc} & a\bar{w}b & \\ \overline{abzx + yy\bar{w}b} & / \quad \backslash & \overline{abzx + yx\bar{w}b} \\ & a\bar{b} \cdot \overline{\bar{x} + \bar{y}} \cdot \bar{w}b & \end{array}$$

Here are some more instances of reciprocity. (There are more.)

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).
 - Show that “almost all” labellings of P are in the image of a certain map Grasp_0 from the matrix space $\mathbb{K}^{p \times (p+q)}$ to $\mathbb{K}^{\hat{P}}$.

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).
 - Show that “almost all” labellings of P are in the image of a certain map Grasp_0 from the matrix space $\mathbb{K}^{P \times (p+q)}$ to $\mathbb{K}^{\hat{P}}$.

Explicitly, if $A \in \mathbb{K}^{P \times (p+q)}$ is any matrix, then $(\text{Grasp}_0 A)(0) = (\text{Grasp}_0 A)(1) = 1$ and

$$(\text{Grasp}_0 A)(i, j) = \frac{\det(A[1 : i \mid i + j - 1 : p + j])}{\det(A[0 : i \mid i + j : p + j])}$$

for all $(i, j) \in P$, where the $A[a : b \mid c : d]$ s are certain submatrices of A . (Note that this map Grasp_0 actually factors through the Grassmannian.)

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).
 - Show that “almost all” labellings of P are in the image of a certain map Grasp_0 from the matrix space $\mathbb{K}^{p \times (p+q)}$ to $\mathbb{K}^{\hat{P}}$.
 - Construct a commutative diagram

$$\begin{array}{ccc} \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \\ \rho \downarrow & & \downarrow R \\ \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \end{array},$$

where ρ is (more or less) rotating the matrix horizontally (last column to front).

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).
 - Show that “almost all” labellings of P are in the image of a certain map Grasp_0 from the matrix space $\mathbb{K}^{p \times (p+q)}$ to $\mathbb{K}^{\hat{P}}$.
 - Construct a commutative diagram

$$\begin{array}{ccc} \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \\ \rho \downarrow & & \downarrow R \\ \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \end{array},$$

where ρ is (more or less) rotating the matrix horizontally (last column to front).

- Conclude that $R^{p+q} = \text{id}$ because $\rho^{p+q} = \text{id}$.

The commutative case

- In 2014, we proved both theorems for commutative \mathbb{K} .
- *Proof outline (inspired by A. Y. Volkov, [arXiv:hep-th/0606094](https://arxiv.org/abs/hep-th/0606094)):*
 - WLOG assume \mathbb{K} is a field (because our claims boil down to polynomial identities).
 - Show that “almost all” labellings of P are in the image of a certain map Grasp_0 from the matrix space $\mathbb{K}^{p \times (p+q)}$ to $\mathbb{K}^{\hat{P}}$.
 - Construct a commutative diagram

$$\begin{array}{ccc} \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \\ \rho \downarrow & & \downarrow R \\ \mathbb{K}^{p \times (p+q)} & \xrightarrow{\text{Grasp}_0} & \mathbb{K}^{\hat{P}} \end{array},$$

where ρ is (more or less) rotating the matrix horizontally (last column to front).

- Conclude that $R^{p+q} = \text{id}$ because $\rho^{p+q} = \text{id}$.
- Reciprocity also easy using Grasp_0 .

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
- Can this be generalized to arbitrary \mathbb{K} ?

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
- Can this be generalized to arbitrary \mathbb{K} ?
- **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
- Can this be generalized to arbitrary \mathbb{K} ?
- **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).
Specifically, redefine Grasp_0 by

$$(\text{Grasp}_0 A)(i, j) = (-1)^i q_{0, i+j-1}^{\{1:i|i+j:p+j\}}(A).$$

The “algebra” works!

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
 - Can this be generalized to arbitrary \mathbb{K} ?
 - **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).
- Specifically, redefine Grasp_0 by

$$(\text{Grasp}_0 A)(i, j) = (-1)^i q_{0, i+j-1}^{\{1:i|i+j:p+j\}}(A).$$

The “algebra” works!

- Unfortunately, the technical parts no longer work:
 - What does “almost all” mean for noncommutative \mathbb{K} ?

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
 - Can this be generalized to arbitrary \mathbb{K} ?
 - **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).
- Specifically, redefine Grasp_0 by

$$(\text{Grasp}_0 A)(i, j) = (-1)^i q_{0, i+j-1}^{\{1:i|i+j:p+j\}}(A).$$

The “algebra” works!

- Unfortunately, the technical parts no longer work:
 - What does “almost all” mean for noncommutative \mathbb{K} ?
 - Can we WLOG assume that \mathbb{K} is a skew field?

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
 - Can this be generalized to arbitrary \mathbb{K} ?
 - **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).
- Specifically, redefine Grasp_0 by

$$(\text{Grasp}_0 A)(i, j) = (-1)^i q_{0, i+j-1}^{\{1:i|i+j:p+j\}}(A).$$

The “algebra” works!

- Unfortunately, the technical parts no longer work:
 - What does “almost all” mean for noncommutative \mathbb{K} ?
 - Can we WLOG assume that \mathbb{K} is a skew field?
No: e.g., the identity $x\overline{y}x = 1$ holds in all skew fields but not in all rings.

First attempts at general proof

- This looks easy; the devil is in the details (particularly the “almost all” part: not just Zariski density but also some rescaling required).
 - Can this be generalized to arbitrary \mathbb{K} ?
 - **In some sense, yes:** Replace determinants by quasideterminants (Gelfand/Retakh, [arXiv:q-alg/9705026](#); see also [arXiv:math/0208146](#)).
- Specifically, redefine Grasp_0 by

$$(\text{Grasp}_0 A)(i, j) = (-1)^i q_{0, i+j-1}^{\{1:i|i+j:p+j\}}(A).$$

The “algebra” works!

- Unfortunately, the technical parts no longer work:
 - What does “almost all” mean for noncommutative \mathbb{K} ?
 - Can we WLOG assume that \mathbb{K} is a skew field?
No: e.g., the identity $x\bar{y}x = 1$ holds in all skew fields but not in all rings.
- We now believe this approach is a dead end.

- New proofs of periodicity and reciprocity in the commutative- \mathbb{K} case were found by Gregg Musiker and Tom Roby in [arXiv:1801.03877](https://arxiv.org/abs/1801.03877).

They proceed by giving an explicit formula for $(R^k f)(i, j)$.
For instance, $(R^3 f)(3, 2)$

$$= \frac{1}{A_{02} + A_{11} + A_{20}} (A_{01}A_{02}A_{11}A_{12} + A_{01}A_{02}A_{12}A_{20} + A_{01}A_{02}A_{20}A_{21} \\ + A_{02}A_{10}A_{12}A_{20} + A_{02}A_{10}A_{20}A_{21} + A_{10}A_{11}A_{20}A_{21}),$$

where

$$A_{ij} := (f(i, j+1) + f(i+1, j)) / f(i+1, j+1).$$

- New proofs of periodicity and reciprocity in the commutative- \mathbb{K} case were found by Gregg Musiker and Tom Roby in [arXiv:1801.03877](https://arxiv.org/abs/1801.03877).

They proceed by giving an explicit formula for $(R^k f)(i, j)$.
For instance, $(R^3 f)(3, 2)$

$$= \frac{1}{A_{02} + A_{11} + A_{20}} (A_{01}A_{02}A_{11}A_{12} + A_{01}A_{02}A_{12}A_{20} + A_{01}A_{02}A_{20}A_{21} \\ + A_{02}A_{10}A_{12}A_{20} + A_{02}A_{10}A_{20}A_{21} + A_{10}A_{11}A_{20}A_{21}),$$

where

$$A_{ij} := (f(i, j+1) + f(i+1, j)) / f(i+1, j+1).$$

- General formula for $(R^k f)(i, j)$ involves sums over NILPs (non-intersecting lattice path families) in numerator and denominator, as well as index shifting and a case split (“small” k and “large” k behave differently).

- New proofs of periodicity and reciprocity in the commutative- \mathbb{K} case were found by Gregg Musiker and Tom Roby in [arXiv:1801.03877](https://arxiv.org/abs/1801.03877).

They proceed by giving an explicit formula for $(R^k f)(i, j)$.
For instance, $(R^3 f)(3, 2)$

$$= \frac{1}{A_{02} + A_{11} + A_{20}} (A_{01}A_{02}A_{11}A_{12} + A_{01}A_{02}A_{12}A_{20} + A_{01}A_{02}A_{20}A_{21} \\ + A_{02}A_{10}A_{12}A_{20} + A_{02}A_{10}A_{20}A_{21} + A_{10}A_{11}A_{20}A_{21}),$$

where

$$A_{ij} := (f(i, j+1) + f(i+1, j)) / f(i+1, j+1).$$

- General formula for $(R^k f)(i, j)$ involves sums over NILPs (non-intersecting lattice path families) in numerator and denominator, as well as index shifting and a case split (“small” k and “large” k behave differently).
- Lattice paths can be generalized to noncommutative \mathbb{K} , but NILPs? Unclear in what order to multiply different paths.

- We are back at square 1: no known theory available.

- We are back at square 1: no known theory available.
- Let's play around with the setting.
Step 1: Introduce notations...

A new beginning

- Fix p, q, P and f . Assume that $R^\ell f$ is well-defined for all necessary ℓ . Let $a = f(0)$ and $b = f(1)$.

A new beginning

- Fix p, q, P and f . Assume that $R^\ell f$ is well-defined for all necessary ℓ . Let $a = f(0)$ and $b = f(1)$.
- For any $x \in \widehat{P}$ and $\ell \in \mathbb{N}$, write

$$x_\ell := (R^\ell f)(x).$$

Thus, $x_0 = f(x)$ and $0_\ell = a$ and $1_\ell = b$.

A new beginning

- Fix p, q, P and f . Assume that $R^\ell f$ is well-defined for all necessary ℓ . Let $a = f(0)$ and $b = f(1)$.
- For any $x \in \widehat{P}$ and $\ell \in \mathbb{N}$, write

$$x_\ell := (R^\ell f)(x).$$

Thus, $x_0 = f(x)$ and $0_\ell = a$ and $1_\ell = b$.

- The definition of R yields

$$(Rf)(v) = \left(\sum_{u < v} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{u > v} \overline{(Rf)(u)}} \quad \text{for each } v \in P.$$

(In both sums, u ranges over \widehat{P} ; this is implied from now on.)

A new beginning

- Fix p, q, P and f . Assume that $R^\ell f$ is well-defined for all necessary ℓ . Let $a = f(0)$ and $b = f(1)$.
- For any $x \in \widehat{P}$ and $\ell \in \mathbb{N}$, write

$$x_\ell := (R^\ell f)(x).$$

Thus, $x_0 = f(x)$ and $0_\ell = a$ and $1_\ell = b$.

- The definition of R yields

$$(Rf)(v) = \left(\sum_{u < v} f(u) \right) \cdot \overline{f(v)} \cdot \overline{\sum_{u > v} (Rf)(u)} \quad \text{for each } v \in P.$$

(In both sums, u ranges over \widehat{P} ; this is implied from now on.)

- In other words,

$$v_1 = \left(\sum_{u < v} u_0 \right) \cdot \overline{v_0} \cdot \overline{\sum_{u > v} u_1} \quad \text{for each } v \in P.$$

- We have just shown that

$$v_1 = \left(\sum_{u < v} u_0 \right) \cdot \overline{v_0} \cdot \overline{\sum_{u > v} u_1} \quad \text{for each } v \in P.$$

- We have just shown that

$$v_1 = \left(\sum_{u < v} u_0 \right) \cdot \overline{v_0} \cdot \overline{\sum_{u > v} u_1} \quad \text{for each } v \in P.$$

- Similarly,

$$v_{\ell+1} = \left(\sum_{u < v} u_\ell \right) \cdot \overline{v_\ell} \cdot \overline{\sum_{u > v} u_{\ell+1}} \quad \text{for each } v \in P \text{ and } \ell \in \mathbb{N}.$$

- We have just shown that

$$v_1 = \left(\sum_{u < v} u_0 \right) \cdot \overline{v_0} \cdot \overline{\sum_{u > v} u_1} \quad \text{for each } v \in P.$$

- Similarly,

$$v_{\ell+1} = \left(\sum_{u < v} u_\ell \right) \cdot \overline{v_\ell} \cdot \overline{\sum_{u > v} u_{\ell+1}} \quad \text{for each } v \in P \text{ and } \ell \in \mathbb{N}.$$

- So far, we have just rewritten our setup using the (more convenient) $x_\ell := (R^\ell f)(x)$ notation.

- We must prove:

periodicity: $x_{p+q} = \bar{a}\bar{b} \cdot x_0 \cdot \bar{a}\bar{b}$;

reciprocity: $x_\ell = a \cdot \overline{y_{\ell-i-j+1}} \cdot b$

if $x = (i, j)$ and $y = (p + 1 - i, q + 1 - j)$.

Simplifying the goal

- We must prove:

periodicity: $x_{p+q} = \overline{ab} \cdot x_0 \cdot \overline{ab}$;

reciprocity: $x_\ell = a \cdot \overline{y_{\ell-i-j+1}} \cdot b$

if $x = (i, j)$ and $y = (p + 1 - i, q + 1 - j)$.

- Periodicity follows from reciprocity: Indeed, if $x = (i, j)$ and $x' = (p + 1 - i, q + 1 - j)$, then

$$\begin{aligned}x_{p+q} &= a \cdot \overline{x'_{p+q-i-j+1}} \cdot b && \text{(by reciprocity)} \\ &= a \cdot \overline{a \cdot \overline{x_0} \cdot b} \cdot b && \text{(by reciprocity again)} \\ &= \overline{ab} \cdot x_0 \cdot \overline{ab}.\end{aligned}$$

Simplifying the goal

- We must prove:

periodicity: $x_{p+q} = \bar{a}\bar{b} \cdot x_0 \cdot \bar{a}b;$

reciprocity: $x_\ell = a \cdot \overline{y_{\ell-i-j+1}} \cdot b$

if $x = (i, j)$ and $y = (p + 1 - i, q + 1 - j)$.

- Periodicity follows from reciprocity: Indeed, if $x = (i, j)$ and $x' = (p + 1 - i, q + 1 - j)$, then

$$\begin{aligned}x_{p+q} &= a \cdot \overline{x'_{p+q-i-j+1}} \cdot b && \text{(by reciprocity)} \\ &= a \cdot \overline{a \cdot \bar{x}_0 \cdot b} \cdot b && \text{(by reciprocity again)} \\ &= \bar{a}\bar{b} \cdot x_0 \cdot \bar{a}b.\end{aligned}$$

Thus, it suffices to prove reciprocity.

Simplifying the goal

- We must prove:

periodicity: $x_{p+q} = \overline{ab} \cdot x_0 \cdot \overline{ab}$;

reciprocity: $x_\ell = a \cdot \overline{y_{\ell-i-j+1}} \cdot b$

if $x = (i, j)$ and $y = (p + 1 - i, q + 1 - j)$.

- Periodicity follows from reciprocity: Indeed, if $x = (i, j)$ and $x' = (p + 1 - i, q + 1 - j)$, then

$$\begin{aligned}x_{p+q} &= a \cdot \overline{x'_{p+q-i-j+1}} \cdot b && \text{(by reciprocity)} \\ &= a \cdot \overline{a \cdot \overline{x_0} \cdot b} \cdot b && \text{(by reciprocity again)} \\ &= \overline{ab} \cdot x_0 \cdot \overline{ab}.\end{aligned}$$

Thus, it suffices to prove reciprocity.

- Moreover, reciprocity in general follows from reciprocity for $\ell = i + j - 1$ (just apply it to $R^k f$ instead of f otherwise).

- A **path** shall mean a sequence $(v_0 \succ v_1 \succ \dots \succ v_k)$ of elements of \widehat{P} . We call it a path from v_0 to v_k .

- A **path** shall mean a sequence $(v_0 \succ v_1 \succ \dots \succ v_k)$ of elements of \widehat{P} . We call it a path from v_0 to v_k .
- For each $v \in P$ and $\ell \in \mathbb{N}$, set

$$A_\ell^v := v_\ell \cdot \overline{\sum_{u < v} u_\ell} \quad \text{and} \quad V_\ell^v := \overline{\sum_{u > v} u_\ell} \cdot \overline{v_\ell}.$$

Also, set $A_\ell^v = V_\ell^v = 1$ when $v \in \{0, 1\}$.

- A **path** shall mean a sequence $(v_0 \succ v_1 \succ \dots \succ v_k)$ of elements of \widehat{P} . We call it a path from v_0 to v_k .
- For each $v \in P$ and $\ell \in \mathbb{N}$, set

$$A_\ell^v := v_\ell \cdot \overline{\sum_{u < v} u_\ell} \quad \text{and} \quad \forall_\ell^v := \overline{\sum_{u > v} u_\ell} \cdot \overline{v_\ell}.$$

Also, set $A_\ell^v = \forall_\ell^v = 1$ when $v \in \{0, 1\}$.

- For any path $\mathbf{p} = (v_0 \succ v_1 \succ \dots \succ v_k)$, set

$$A_\ell^{\mathbf{p}} := A_\ell^{v_0} A_\ell^{v_1} \dots A_\ell^{v_k} \quad \text{and} \quad \forall_\ell^{\mathbf{p}} := \forall_\ell^{v_0} \forall_\ell^{v_1} \dots \forall_\ell^{v_k}.$$

- A **path** shall mean a sequence $(v_0 \succ v_1 \succ \dots \succ v_k)$ of elements of \widehat{P} . We call it a path from v_0 to v_k .
- For each $v \in P$ and $\ell \in \mathbb{N}$, set

$$A_\ell^v := v_\ell \cdot \overline{\sum_{u < v} u_\ell} \quad \text{and} \quad \forall_\ell^v := \overline{\sum_{u > v} u_\ell} \cdot \overline{v_\ell}.$$

Also, set $A_\ell^v = \forall_\ell^v = 1$ when $v \in \{0, 1\}$.

- For any path $\mathbf{p} = (v_0 \succ v_1 \succ \dots \succ v_k)$, set

$$A_\ell^{\mathbf{p}} := A_\ell^{v_0} A_\ell^{v_1} \dots A_\ell^{v_k} \quad \text{and} \quad \forall_\ell^{\mathbf{p}} := \forall_\ell^{v_0} \forall_\ell^{v_1} \dots \forall_\ell^{v_k}.$$

- If u and v are elements of \widehat{P} , set

$$A_\ell^{u \rightarrow v} := \sum_{\mathbf{p} \text{ is a path from } u \text{ to } v} A_\ell^{\mathbf{p}} \quad \text{and}$$

$$\forall_\ell^{u \rightarrow v} := \sum_{\mathbf{p} \text{ is a path from } u \text{ to } v} \forall_\ell^{\mathbf{p}}.$$

- **Path formulas:**

(a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

(b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* The ℓ is constant. Hence, we omit it, writing \forall^ν for \forall_ℓ^ν .

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* The ℓ is constant. Hence, we omit it, writing \forall^ν for \forall_ℓ^ν .

- (a) Rewrite the claim as $\forall^{1 \rightarrow u} = b\overline{u}_\ell$.

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* The ℓ is constant. Hence, we omit it, writing \forall^ν for \forall_ℓ^ν .

- (a) Rewrite the claim as $\forall^{1 \rightarrow u} = b\overline{u}_\ell$.

- Prove this by downwards induction on u .

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\mathcal{V}_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* The ℓ is constant. Hence, we omit it, writing \mathcal{V}^ν for \mathcal{V}_ℓ^ν .

- (a) Rewrite the claim as $\mathcal{V}^{1 \rightarrow u} = b\overline{u}_\ell$.

Prove this by downwards induction on u .

Induction step: Given $v \in P$ such that $\mathcal{V}^{1 \rightarrow u} = b\overline{u}_\ell$ for all $u \succ v$. Since any path $1 \rightarrow v$ passes through a unique $u \succ v$, we have

$$\begin{aligned} \mathcal{V}^{1 \rightarrow v} &= \sum_{u \succ v} \mathcal{V}^{1 \rightarrow u} \mathcal{V}^v = \sum_{u \succ v} b\overline{u}_\ell \mathcal{V}^v && \text{(by induction hypothesis)} \\ &= b\overline{v}_\ell && \text{(by definition of } \mathcal{V}^v \text{),} \quad \text{qed.} \end{aligned}$$

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* The ℓ is constant. Hence, we omit it, writing \forall^ν for \forall_ℓ^ν .

- (b) Analogous, but use upwards induction instead.

- **Path formulas:**

(a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

(b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

(c) We have

$$u_\ell = \overline{\forall_\ell^{(p,q) \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

(d) We have

$$u_\ell = A_\ell^{u \rightarrow (1,1)} \cdot a \quad \text{for each } u \in P.$$

- **Path formulas:**

- (a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

- (c) We have

$$u_\ell = \overline{\forall_\ell^{(p,q) \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

- (d) We have

$$u_\ell = A_\ell^{u \rightarrow (1,1)} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* Each path $1 \rightarrow u$ begins with the step $1 \succ (p, q)$. Thus, $\forall_\ell^{1 \rightarrow u} = \forall_\ell^{(p,q) \rightarrow u}$ (since $\forall_\ell^1 = 1$). Hence, (c) follows from (a).

- **Path formulas:**

(a) We have

$$u_\ell = \overline{\forall_\ell^{1 \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

(b) We have

$$u_\ell = A_\ell^{u \rightarrow 0} \cdot a \quad \text{for each } u \in P.$$

(c) We have

$$u_\ell = \overline{\forall_\ell^{(p,q) \rightarrow u}} \cdot b \quad \text{for each } u \in P.$$

(d) We have

$$u_\ell = A_\ell^{u \rightarrow (1,1)} \cdot a \quad \text{for each } u \in P.$$

- *Proof idea:* Each path $1 \rightarrow u$ begins with the step $1 \succ (p, q)$.

Thus, $\forall_\ell^{1 \rightarrow u} = \forall_\ell^{(p,q) \rightarrow u}$ (since $\forall_\ell^1 = 1$). Hence, (c) follows from (a).

Similarly, (d) follows from (b).

- **Transition equation in A - \mathcal{V} -form:**

$$\mathcal{V}_{\ell+1}^v = A_\ell^v \quad \text{for each } v \in \widehat{P} \text{ and } \ell \in \mathbb{N}.$$

- **Transition equation in A - \forall -form:**

$$\forall_{\ell+1}^v = A_\ell^v \quad \text{for each } v \in \widehat{P} \text{ and } \ell \in \mathbb{N}.$$

- *Proof idea:* Above we showed that

$$v_{\ell+1} = \left(\sum_{u \leq v} u_\ell \right) \cdot \overline{v_\ell} \cdot \overline{\sum_{u \geq v} \overline{u_{\ell+1}}}.$$

Take reciprocals on both sides, multiply by $\overline{\sum_{u \geq v} \overline{u_{\ell+1}}}$ and rewrite using $\forall_{\ell+1}^v$ and A_ℓ^v .

- **Transition equation in A - \forall -form:**

$$\forall_{\ell+1}^v = A_\ell^v \quad \text{for each } v \in \widehat{P} \text{ and } \ell \in \mathbb{N}.$$

- *Proof idea:* Above we showed that

$$v_{\ell+1} = \left(\sum_{u \leq v} u_\ell \right) \cdot \overline{v_\ell} \cdot \overline{\sum_{u \geq v} \overline{u_{\ell+1}}}.$$

Take reciprocals on both sides, multiply by $\overline{\sum_{u \geq v} \overline{u_{\ell+1}}}$ and rewrite using $\forall_{\ell+1}^v$ and A_ℓ^v .

- As a consequence of $\forall_{\ell+1}^v = A_\ell^v$, we have

$$\forall_{\ell+1}^{\mathbf{p}} = A_\ell^{\mathbf{p}} \quad \text{for each path } \mathbf{p} \text{ and each } \ell \in \mathbb{N}.$$

- **Transition equation in A - \forall -form:**

$$\forall_{\ell+1}^v = A_\ell^v \quad \text{for each } v \in \widehat{P} \text{ and } \ell \in \mathbb{N}.$$

- *Proof idea:* Above we showed that

$$v_{\ell+1} = \left(\sum_{u \leq v} u_\ell \right) \cdot \overline{v_\ell} \cdot \overline{\sum_{u \geq v} \overline{u_{\ell+1}}}.$$

Take reciprocals on both sides, multiply by $\overline{\sum_{u \geq v} \overline{u_{\ell+1}}}$ and rewrite using $\forall_{\ell+1}^v$ and A_ℓ^v .

- As a consequence of $\forall_{\ell+1}^v = A_\ell^v$, we have

$$\forall_{\ell+1}^{\mathbf{p}} = A_\ell^{\mathbf{p}} \quad \text{for each path } \mathbf{p} \text{ and each } \ell \in \mathbb{N}.$$

Hence, $\forall_{\ell+1}^{u \rightarrow v} = A_\ell^{u \rightarrow v}$ for any $u, v \in \widehat{P}$.

- Now, for the bottommost element $(1, 1)$ of P , we have

$$\begin{aligned}(1, 1)_1 &= \overline{V_1^{(p,q) \rightarrow (1,1)}} \cdot b && \text{(by path formula (c))} \\ &= \overline{A_0^{(p,q) \rightarrow (1,1)}} \cdot b && \text{(since } V_{\ell+1}^{u \rightarrow v} = A_{\ell}^{u \rightarrow v} \text{)} \\ &= a \cdot \overline{(p, q)_0} \cdot b && \text{(by path formula (d)).}\end{aligned}$$

Thus, reciprocity is proved for $i = j = 1$.

- Now, for the bottommost element $(1, 1)$ of P , we have


$$\begin{aligned}(1, 1)_1 &= \overline{V_1^{(p,q) \rightarrow (1,1)}} \cdot b && \text{(by path formula (c))} \\ &= \overline{A_0^{(p,q) \rightarrow (1,1)}} \cdot b && \text{(since } V_{\ell+1}^{u \rightarrow v} = A_\ell^{u \rightarrow v}\text{)} \\ &= a \cdot \overline{(p, q)_0} \cdot b && \text{(by path formula (d)).}\end{aligned}$$

Thus, reciprocity is proved for $i = j = 1$.

- What now?

The case $j = 1$ suffices: part 1

- We can simplify our goal one bit further. Consider the “neighborhood” of an element of our rectangle P :




(where the **rank** of an $(i, j) \in P$ is defined to be $i + j - 1$). Say we have shown (our “induction hypotheses”) that reciprocity holds for each of s, t, m, u ; that is, we have

$$\begin{aligned}
 s_\ell &= a \cdot \overline{s'_{\ell-(k-1)}} \cdot b, & t_\ell &= a \cdot \overline{t'_{\ell-(k-1)}} \cdot b, \\
 m_\ell &= a \cdot \overline{m'_{\ell-k}} \cdot b, & u_\ell &= a \cdot \overline{u'_{\ell-(k+1)}} \cdot b
 \end{aligned}$$

for all sufficiently high ℓ , where x' denotes the antipode of x (that is, if $x = (i, j)$, then $x' = (p + 1 - i, q + 1 - j)$).

The case $j = 1$ suffices: part 1

- We can simplify our goal one bit further. Consider the “neighborhood” of an element of our rectangle P :



(where the **rank** of an $(i, j) \in P$ is defined to be $i + j - 1$). Say we have shown (our “induction hypotheses”) that reciprocity holds for each of s, t, m, u ; that is, we have

$$\begin{aligned}
 s_\ell &= a \cdot \overline{s'_{\ell-(k-1)}} \cdot b, & t_\ell &= a \cdot \overline{t'_{\ell-(k-1)}} \cdot b, \\
 m_\ell &= a \cdot \overline{m'_{\ell-k}} \cdot b, & u_\ell &= a \cdot \overline{u'_{\ell-(k+1)}} \cdot b
 \end{aligned}$$

for all sufficiently high ℓ , where x' denotes the antipode of x (that is, if $x = (i, j)$, then $x' = (p + 1 - i, q + 1 - j)$).

Claim: Then, reciprocity also holds for v ; that is, we have $v_\ell = a \cdot \overline{v'_{\ell-(k+1)}} \cdot b$ for all $\ell \geq k + 1$.

The case $j = 1$ suffices: part 2

- *Proof idea.* Fix $\ell \geq k + 1$, and compare the transition equations

$$m_\ell = (s_{\ell-1} + t_{\ell-1}) \cdot \overline{m_{\ell-1}} \cdot \overline{u_\ell + v_\ell} \quad \text{and}$$

$$m'_{\ell-k} = (u'_{\ell-k-1} + v'_{\ell-k-1}) \cdot \overline{m'_{\ell-k-1}} \cdot \overline{s'_{\ell-k} + t'_{\ell-k}}$$

using the induction hypotheses $m_\ell = a \cdot \overline{m'_{\ell-k}} \cdot b$,


$$s_{\ell-1} = a \cdot \overline{s'_{\ell-k}} \cdot b,$$

$$t_{\ell-1} = a \cdot \overline{t'_{\ell-k}} \cdot b,$$

$$m_{\ell-1} = a \cdot \overline{m'_{\ell-1-k}} \cdot b,$$

$$u_\ell = a \cdot \overline{u'_{\ell-(k+1)}} \cdot b,$$

noting that



The case $j = 1$ suffices: part 2

- *Proof idea.* Fix $\ell \geq k + 1$, and compare the transition equations

$$m_\ell = (s_{\ell-1} + t_{\ell-1}) \cdot \overline{m_{\ell-1}} \cdot \overline{u_\ell + v_\ell} \quad \text{and}$$

$$m'_{\ell-k} = (u'_{\ell-k-1} + v'_{\ell-k-1}) \cdot \overline{m'_{\ell-k-1}} \cdot \overline{s'_{\ell-k} + t'_{\ell-k}}$$

using the induction hypotheses $m_\ell = a \cdot \overline{m'_{\ell-k}} \cdot b$,

$$s_{\ell-1} = a \cdot \overline{s'_{\ell-k}} \cdot b, \quad t_{\ell-1} = a \cdot \overline{t'_{\ell-k}} \cdot b,$$

$$m_{\ell-1} = a \cdot \overline{m'_{\ell-1-k}} \cdot b, \quad u_\ell = a \cdot \overline{u'_{\ell-(k+1)}} \cdot b,$$

noting that

$$\begin{array}{ccc}
 u & & v \\
 & \diagdown & / \\
 & m & \\
 & / & \diagdown \\
 s & & t
 \end{array}
 \implies
 \begin{array}{ccc}
 t' & & s' \\
 & \diagdown & / \\
 & m' & \\
 & / & \diagdown \\
 v' & & u'
 \end{array}
 .$$

After subtracting $u_\ell = a \cdot \overline{u'_{\ell-(k+1)}} \cdot b$, out comes

$$v_\ell = a \cdot \overline{v'_{\ell-(k+1)}} \cdot b.$$

The case $j = 1$ suffices: part 2

- *Proof idea.* Fix $\ell \geq k + 1$, and compare the transition equations

$$m_\ell = (s_{\ell-1} + t_{\ell-1}) \cdot \overline{m_{\ell-1}} \cdot \overline{u_\ell + v_\ell} \quad \text{and}$$

$$m'_{\ell-k} = (u'_{\ell-k-1} + v'_{\ell-k-1}) \cdot \overline{m'_{\ell-k-1}} \cdot \overline{s'_{\ell-k} + t'_{\ell-k}}$$

using the induction hypotheses $m_\ell = a \cdot \overline{m'_{\ell-k}} \cdot b$,


$$s_{\ell-1} = a \cdot \overline{s'_{\ell-k}} \cdot b,$$

$$t_{\ell-1} = a \cdot \overline{t'_{\ell-k}} \cdot b,$$

$$m_{\ell-1} = a \cdot \overline{m'_{\ell-1-k}} \cdot b,$$

$$u_\ell = a \cdot \overline{u'_{\ell-(k+1)}} \cdot b,$$

noting that



- This argument still works if s , t or u does not exist.

The case $j = 1$ suffices: part 2

- *Proof idea.* Fix $\ell \geq k + 1$, and compare the transition equations

$$m_\ell = (s_{\ell-1} + t_{\ell-1}) \cdot \overline{m_{\ell-1}} \cdot \overline{u_\ell + v_\ell} \quad \text{and}$$

$$m'_{\ell-k} = (u'_{\ell-k-1} + v'_{\ell-k-1}) \cdot \overline{m'_{\ell-k-1}} \cdot \overline{s'_{\ell-k} + t'_{\ell-k}}$$

using the induction hypotheses $m_\ell = a \cdot \overline{m'_{\ell-k}} \cdot b$,


$$s_{\ell-1} = a \cdot \overline{s'_{\ell-k}} \cdot b,$$

$$t_{\ell-1} = a \cdot \overline{t'_{\ell-k}} \cdot b,$$

$$m_{\ell-1} = a \cdot \overline{m'_{\ell-1-k}} \cdot b,$$

$$u_\ell = a \cdot \overline{u'_{\ell-(k+1)}} \cdot b,$$

noting that



- This argument still works if s , t or u does not exist.
- Thus, in order to prove reciprocity for all (i, j) , it suffices (by induction) to prove it in the case when $j = 1$.

Where are we?

- So we have proved reciprocity for $i = j = 1$, and we need to prove it for $j = 1$.

Where are we?

- So we have proved reciprocity for $i = j = 1$, and we need to prove it for $j = 1$.
- The next case to try is $(i, j) = (2, 1)$. We need to show that

$$(2, 1)_2 = a \cdot \overline{(p-1, q)_0} \cdot b.$$

Where are we?

- So we have proved reciprocity for $i = j = 1$, and we need to prove it for $j = 1$.
- The next case to try is $(i, j) = (2, 1)$. We need to show that

$$(2, 1)_2 = a \cdot \overline{(p-1, q)_0} \cdot b.$$

- Using the path formulas (as in the case $i = j = 1$), we can boil this down to

$$A_1^{(p,q) \rightarrow (2,1)} = V_1^{(p-1,q) \rightarrow (1,1)}.$$

Where are we?

- So we have proved reciprocity for $i = j = 1$, and we need to prove it for $j = 1$.
- The next case to try is $(i, j) = (2, 1)$. We need to show that

$$(2, 1)_2 = a \cdot \overline{(p-1, q)_0} \cdot b.$$

- Using the path formulas (as in the case $i = j = 1$), we can boil this down to

$$A_1^{(p,q) \rightarrow (2,1)} = V_1^{(p-1,q) \rightarrow (1,1)}.$$

Note the lack of rowmotion in this formula! The ℓ here is constantly 1, so it is a property of a single labeling. Thus, we drop the subscripts.

- **Our new goal:** Prove that

$$A^{(p,q) \rightarrow (2,1)} = V^{(p-1,q) \rightarrow (1,1)}.$$


The conversion lemma

- More generally:
- **Conversion lemma:**
Let u and u' be two adjacent elements on the top-right edge of P (that is, $u = (k, q)$ and $u' = (k - 1, q)$). Let d and d' be two adjacent elements on the bottom-left edge of P (that is, $d = (i, 1)$ and $d' = (i - 1, 1)$). Then,

$$A_\ell^{u \rightarrow d} = \forall_\ell^{u' \rightarrow d'} \quad \text{for each } \ell \in \mathbb{N}.$$

In short:

$$A^{u \rightarrow d} = \forall^{u' \rightarrow d'}.$$



- If we can prove the conversion lemma, we will obtain reciprocity not only for $(i, j) = (2, 1)$, but also for all (i, j) on the bottom-left edge of P (that is, for the entire case $j = 1$), because we can argue as follows:

$$\begin{aligned}
 (i, 1)_i &= \overline{\mathcal{V}_i^{(p,q) \rightarrow (i,1)}} \cdot b \\
 &= \overline{A_{i-1}^{(p,q) \rightarrow (i,1)}} \cdot b \\
 &= \overline{\mathcal{V}_{i-1}^{(p-1,q) \rightarrow (i-1,1)}} \cdot b \\
 &= \overline{A_{i-2}^{(p-1,q) \rightarrow (i-1,1)}} \cdot b \\
 &= \overline{\mathcal{V}_{i-2}^{(p-2,q) \rightarrow (i-2,1)}} \cdot b \\
 &= \dots \\
 &= \overline{\mathcal{V}_1^{(p-i+1,q) \rightarrow (1,1)}} \cdot b \\
 &= \overline{A_0^{(p-i+1,q) \rightarrow (1,1)}} \cdot b \\
 &= a \cdot \overline{(p-i+1, q)_0} \cdot b
 \end{aligned}$$

(by path formula **(c)**)

(since $\mathcal{V}_{\ell+1}^{u \rightarrow v} = A_{\ell}^{u \rightarrow v}$)

(by the conversion lemma)

(since $\mathcal{V}_{\ell+1}^{u \rightarrow v} = A_{\ell}^{u \rightarrow v}$)

(by the conversion lemma)

(by the conversion lemma)

(since $\mathcal{V}_{\ell+1}^{u \rightarrow v} = A_{\ell}^{u \rightarrow v}$)

(by path formula **(d)**).

- This proves reciprocity

$$(i, 1)_\ell = a \cdot \overline{(p - i + 1, q)_{\ell-i}} \cdot b$$

for $\ell = i$.

- This proves reciprocity

$$(i, 1)_\ell = a \cdot \overline{(p - i + 1, q)_{\ell-i}} \cdot b$$

for $\ell = i$.

The case $\ell > i$ follows by applying this to $R^{\ell-i}f$ instead of f .

- This proves reciprocity

$$(i, 1)_\ell = a \cdot \overline{(p - i + 1, q)_{\ell-i}} \cdot b$$


for $\ell = i$.

The case $\ell > i$ follows by applying this to $R^{\ell-i}f$ instead of f .

- Thus, we only need to prove the conversion lemma. We can now drop all subscripts forever!

Proving the conversion lemma: the intuition


- Let us again look at the picture:



We must prove $A^{u \rightarrow d} = \forall u' \rightarrow d'$.

Proving the conversion lemma: the intuition

- Let us again look at the picture:



We must prove $A^{u \rightarrow d} = \forall u' \rightarrow d'$.

- How do we interpolate between paths $u \rightarrow d$ and paths $u' \rightarrow d'$?

- We define a **path-jump-path** to be a sequence

$$\mathbf{p} = (v_0 \succ v_1 \succ \cdots \succ v_i \blacktriangleright v_{i+1} \succ v_{i+2} \succ \cdots \succ v_k)$$

of elements of P , where the relation $x \blacktriangleright y$ means “ y is one step down and some steps to the right of x ” (that is, if $x = (r, s)$, then $y = (r - k, s + k - 1)$ for some $k > 0$).

We say that this path-jump-path \mathbf{p} has **jump at i** .

Proving the conversion lemma: path-jump-paths


- We define a **path-jump-path** to be a sequence

$$\mathbf{p} = (v_0 \succ v_1 \succ \cdots \succ v_i \blacktriangleright v_{i+1} \succ v_{i+2} \succ \cdots \succ v_k)$$

of elements of P , where the relation $x \blacktriangleright y$ means “ y is one step down and some steps to the right of x ” (that is, if $x = (r, s)$, then $y = (r - k, s + k - 1)$ for some $k > 0$).

We say that this path-jump-path \mathbf{p} has **jump at i** .

Example of a path-jump-path:



(The red edge is the jump.)

- We define a **path-jump-path** to be a sequence

$$\mathbf{p} = (v_0 \succ v_1 \succ \cdots \succ v_i \blacktriangleright v_{i+1} \succ v_{i+2} \succ \cdots \succ v_k)$$

of elements of P , where the relation $x \blacktriangleright y$ means “ y is one step down and some steps to the right of x ” (that is, if $x = (r, s)$, then $y = (r - k, s + k - 1)$ for some $k > 0$).

We say that this path-jump-path \mathbf{p} has **jump at i** .

For any such path-jump-path \mathbf{p} , we set

$$E_{\mathbf{p}} := A^{v_0} A^{v_1} \cdots A^{v_{i-1}} v_i \overline{v_{i+1}} \mathcal{V}^{v_{i+2}} \mathcal{V}^{v_{i+3}} \cdots \mathcal{V}^{v_k}.$$

(Here, we are omitting the ℓ subscripts – so v_i means $(v_i)_{\ell}$ and v_{i+1} means $(v_{i+1})_{\ell}$.)

Proving the conversion lemma: path-jump-paths

- We define a **path-jump-path** to be a sequence

$$\mathbf{p} = (v_0 \succ v_1 \succ \cdots \succ v_i \blacktriangleright v_{i+1} \succ v_{i+2} \succ \cdots \succ v_k)$$

of elements of P , where the relation $x \blacktriangleright y$ means “ y is one step down and some steps to the right of x ” (that is, if $x = (r, s)$, then $y = (r - k, s + k - 1)$ for some $k > 0$).

We say that this path-jump-path \mathbf{p} has **jump at i** .

For any such path-jump-path \mathbf{p} , we set

$$E_{\mathbf{p}} := A^{v_0} A^{v_1} \cdots A^{v_{i-1}} v_i \overline{v_{i+1}} \forall^{v_{i+2}} \forall^{v_{i+3}} \cdots \forall^{v_k}.$$

- Now, if $k = \text{rank } u - \text{rank}(d')$, then

$$A^{u \rightarrow d} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } k-1}} E_{\mathbf{p}},$$

since $A^d = d \overline{d'}$, and similarly

$$\forall^{u' \rightarrow d'} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } 0}} E_{\mathbf{p}}.$$

- So we need to show that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } k-1}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } 0}} E_{\mathbf{p}}.$$

- So we need to show that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } k-1}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } 0}} E_{\mathbf{p}}.$$

- Reasonable to expect that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i+1}} E_{\mathbf{p}}$$

for each $0 \leq i < k - 1$.

- So we need to show that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } k-1}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } 0}} E_{\mathbf{p}}.$$

- Reasonable to expect that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i+1}} E_{\mathbf{p}}$$

for each $0 \leq i < k - 1$.

- This is indeed true and can be proved by a “local” argument (rewriting two consecutive steps of the path).

- So we need to show that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } k-1}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } 0}} E_{\mathbf{p}}.$$

- Reasonable to expect that

$$\sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i}} E_{\mathbf{p}} = \sum_{\substack{\mathbf{p} \text{ is a path-jump-path } u \rightarrow d' \\ \text{with jump at } i+1}} E_{\mathbf{p}}$$

for each $0 \leq i < k - 1$.

- This is indeed true and can be proved by a “local” argument (rewriting two consecutive steps of the path).
- This is similar to the “zipper argument” in lattice models. (Is there a Yang–Baxter equation lurking?)

- Modulo the details omitted, this finishes the proof of the reciprocity theorem.

Proving the conversion lemma: the civilized version, part 1

- Modulo the details omitted, this finishes the proof of the reciprocity theorem.
- However, the path-jump-path argument is somewhat messy. We can make it slicker by rewriting it in matrix notation:

Proving the conversion lemma: the civilized version, part 1

- Modulo the details omitted, this finishes the proof of the reciprocity theorem.
- However, the path-jump-path argument is somewhat messy. We can make it slicker by rewriting it in matrix notation:
- Define three $P \times P$ -matrices \mathbf{A} , \mathbf{V} and \mathbf{U} by

$$\begin{aligned}\mathbf{A}_{x,y} &:= A^x [x \succcurlyeq y], & \mathbf{V}_{x,y} &:= V^y [x \succcurlyeq y], \\ \mathbf{U}_{x,y} &:= x\bar{y} [x \blacktriangleright y] & & \text{for all } x, y \in P.\end{aligned}$$

Here, $[\mathcal{A}]$ is the Iverson bracket (i.e., truth value) of a statement \mathcal{A} ; the relation $x \blacktriangleright y$ means “ y is one step down and some steps to the right of x ” as before. And again, we are omitting the ℓ subscripts, so $x\bar{y}$ actually means $x_\ell \bar{y}_\ell$.


- Now, we claim that

$$\mathbf{AU} = \mathbf{UV}.$$

Proving the conversion lemma: the civilized version, part 2

- Now, we claim that $\mathbf{AU} = \mathbf{UV}$.

Indeed, this follows easily from the following neat lemma: If



are four adjacent elements of P , then


$$\bar{w} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot v \quad \text{and} \quad \bar{v} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot w.$$

(The u and d here are unrelated to the u and d from the conversion lemma!)

Proving the conversion lemma: the civilized version, part 2

- Now, we claim that $\mathbf{AU} = \mathbf{UV}$.

Indeed, this follows easily from the following neat lemma: If



are four adjacent elements of P , then

$$\bar{w} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot v \quad \text{and} \quad \bar{v} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot w.$$

- From $\mathbf{AU} = \mathbf{UV}$, we easily obtain


$$\mathbf{A}^{\circ k} \mathbf{U} = \mathbf{UV}^{\circ k} \quad \text{for any } k \in \mathbb{N},$$

where $A^{\circ k}$ means the k -th power of a matrix A .

Proving the conversion lemma: the civilized version, part 2

- Now, we claim that $\mathbf{AU} = \mathbf{UV}$.

Indeed, this follows easily from the following neat lemma: If



are four adjacent elements of P , then

$$\bar{w} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot v \quad \text{and} \quad \bar{v} \cdot \nabla^d \cdot d = \bar{u} \cdot A^u \cdot w.$$

- From $\mathbf{AU} = \mathbf{UV}$, we easily obtain

$$\mathbf{A}^{\circ k} \mathbf{U} = \mathbf{UV}^{\circ k} \quad \text{for any } k \in \mathbb{N},$$

where $A^{\circ k}$ means the k -th power of a matrix A .

- Setting $k = \text{rank } u - \text{rank } d$ and comparing the (u, d') -entries of both sides, we quickly obtain $A^{u \rightarrow d} = \nabla^{u' \rightarrow d'}$ (since $x \blacktriangleright d'$ holds only for $x = d$, and since $u \blacktriangleright x$ holds only for $x = u'$). This proves the conversion lemma again.

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.

Is that all? Part 1: Semirings

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.
- However, in some sense they are still imperfect.

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.
- However, in some sense they are still imperfect.
- **Recall:** Classical rowmotion is (a restriction of) birational rowmotion on the tropical **semifield**.

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.
- However, in some sense they are still imperfect.
- **Recall:** Classical rowmotion is (a restriction of) birational rowmotion on the tropical **semifield**. Semifields are not rings! (No subtraction.)

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.
- However, in some sense they are still imperfect.
- **Recall:** Classical rowmotion is (a restriction of) birational rowmotion on the tropical **semifield**.
Semifields are not rings! (No subtraction.)
In the **commutative** case, the theorems hold for semifields (and, more generally, commutative semirings) because they hold for fields and because they are “essentially” polynomial identities (once you clear denominators).

- We consider these new proofs to be simpler and nicer than our 2014 one for the commutative case.
- However, in some sense they are still imperfect.
- **Recall:** Classical rowmotion is (a restriction of) birational rowmotion on the tropical **semifield**.

Semifields are not rings! (No subtraction.)

In the **commutative** case, the theorems hold for semifields (and, more generally, commutative semirings) because they hold for fields and because they are “essentially” polynomial identities (once you clear denominators).

This **fails** for noncommutative \mathbb{K} !

- **Scary example** ([David Speyer, MathOverflow #401273](#)): If x and y are two elements of a ring such that $x + y$ is invertible, then

$$x \cdot \overline{x + y} \cdot y = y \cdot \overline{x + y} \cdot x.$$

But this is not true if “ring” is replaced by “semiring”!

- Thus, we are left with a

Question:

Are the periodicity and reciprocity theorems still true if “ring” is replaced by “semiring”? (I.e., we no longer require \mathbb{K} to have a subtraction.)

- Thus, we are left with a

Question:

Are the periodicity and reciprocity theorems still true if “ring” is replaced by “semiring”? (I.e., we no longer require \mathbb{K} to have a subtraction.)

- Note that the main hurdle is the argument that reduced the general case to the $j = 1$ case. That argument used subtraction!

- Thus, we are left with a

Question:

Are the periodicity and reciprocity theorems still true if “ring” is replaced by “semiring”? (I.e., we no longer require \mathbb{K} to have a subtraction.)

- Note that the main hurdle is the argument that reduced the general case to the $j = 1$ case. That argument used subtraction!
- We have partial results, e.g., for $p = q = 3$ and for $p = 2$.


- Other posets remain to be studied.

Conjecture:

Let P be the triangle-shaped poset $\Delta(p)$ or its reflection $\nabla(p)$. Let $f \in \mathbb{K}^{\hat{P}}$ be a labelling such that $R^P f$ exists. Let $a = f(0)$ and $b = f(1)$. Then, for each $x \in \hat{P}$, we have

$$(R^P f)(x) = a\bar{b} \cdot f(x') \cdot \bar{a}b,$$

where x' is the reflection of x across the y -axis.



- Other posets remain to be studied.

Conjecture:

Let P be the triangle-shaped poset $\Delta(p)$ or its reflection $\nabla(p)$. Let $f \in \mathbb{K}^{\hat{P}}$ be a labelling such that $R^P f$ exists. Let $a = f(0)$ and $b = f(1)$. Then, for each $x \in \hat{P}$, we have

$$(R^P f)(x) = a\bar{b} \cdot f(x') \cdot \bar{a}b,$$

where x' is the reflection of x across the y -axis.

- We have a similar conjecture for other kinds of triangles and (still unproved even in the commutative case!) for trapezoids.

- Other posets remain to be studied.


Conjecture:

Let P be the triangle-shaped poset $\Delta(p)$ or its reflection $\nabla(p)$. Let $f \in \mathbb{K}^{\hat{P}}$ be a labelling such that $R^P f$ exists. Let $a = f(0)$ and $b = f(1)$. Then, for each $x \in \hat{P}$, we have

$$(R^P f)(x) = a\bar{b} \cdot f(x') \cdot \bar{a}b,$$

where x' is the reflection of x across the y -axis.

- We have a similar conjecture for other kinds of triangles and (still unproved even in the commutative case!) for trapezoids.
- As already mentioned, other simple posets such as



do not have periodic behavior for noncommutative \mathbb{K} .

- Other posets remain to be studied.

Conjecture:

Let P be the triangle-shaped poset $\Delta(p)$ or its reflection $\nabla(p)$. Let $f \in \mathbb{K}^{\hat{P}}$ be a labelling such that $R^P f$ exists. Let $a = f(0)$ and $b = f(1)$. Then, for each $x \in \hat{P}$, we have

$$(R^P f)(x) = a\bar{b} \cdot f(x') \cdot \bar{a}b,$$

where x' is the reflection of x across the y -axis.

- We have a similar conjecture for other kinds of triangles and (still unproved even in the commutative case!) for trapezoids.

Question:

What other results like ours are known in the noncommutative case?

- **Tom Roby**: collaboration
- **Mathematisches Forschungsinstitut Oberwolfach**: hospitality in July/August 2021
- **Banff International Research Station**: 2021 conference where this was first presented
- **Svante Linusson**: invitation
- **Michael Joseph, Tim Champion, Max Glick, Maxim Kontsevich, Gregg Musiker, Pace Nielsen, James Propp, Pasha Pylyavskyy, Bruce Sagan, Roland Speicher, David Speyer, Hugh Thomas, and Jurij Volcic**: discussions
- **Sage and Sage-combinat**: computations
- **the birational combinatorics community**: keeping the subject exciting since 2013
- **you**: your patience

Some references

- David Einstein, James Propp, *Combinatorial, piecewise-linear, and birational homomesy for products of two chains*, 2013.
<http://arxiv.org/abs/1310.5294>
- David Einstein, James Propp, *Piecewise-linear and birational toggling*, 2014. <https://arxiv.org/abs/1404.3455>
- Darij Grinberg, Tom Roby, *Iterative properties of birational rowmotion*, 2014. <http://arxiv.org/abs/1402.6178>
- Michael Joseph, Tom Roby, *Birational and noncommutative lifts of antichain toggling and rowmotion*, 2019.
<https://arxiv.org/abs/1909.09658>
- Michael Joseph, Tom Roby, *A birational lifting of the Stanley-Thomas word on products of two chains*, 2020.
<https://arxiv.org/abs/2001.03811>
- Gregg Musiker, Tom Roby, *Paths to Understanding Birational Rowmotion on Products of Two Chains*, 2019.
<https://arxiv.org/abs/1801.03877>

- **Zamolodchikov periodicity conjecture in type AA** (proved by A. Yu. Volkov, [arXiv:hep-th/0606094v1](https://arxiv.org/abs/hep-th/0606094v1)): Let r and s be positive integers. Let $Y_{i,j,k}$ be elements of a commutative ring for $i \in [r]$ and $j \in [s]$ and $k \in \mathbb{Z}$. Assume that

$$Y_{i,j,k+1} Y_{i,j,k-1} = \frac{(1 + Y_{i+1,j,k})(1 + Y_{i-1,j,k})}{(1 + 1/Y_{i,j+1,k})(1 + 1/Y_{i,j-1,k})}$$

for all i, j, k , where sums involving “off-grid” points (e.g., $1 + Y_{0,j,k}$) are understood as 1.

Then, $Y_{i,j,k+2(r+s+2)} = Y_{i,j,k}$ for all i, j, k .

- **Zamolodchikov periodicity conjecture in type AA** (proved by A. Yu. Volkov, [arXiv:hep-th/0606094v1](https://arxiv.org/abs/hep-th/0606094v1)): Let r and s be positive integers. Let $Y_{i,j,k}$ be elements of a commutative ring for $i \in [r]$ and $j \in [s]$ and $k \in \mathbb{Z}$. Assume that

$$Y_{i,j,k+1} Y_{i,j,k-1} = \frac{(1 + Y_{i+1,j,k})(1 + Y_{i-1,j,k})}{(1 + 1/Y_{i,j+1,k})(1 + 1/Y_{i,j-1,k})}$$

for all i, j, k , where sums involving “off-grid” points (e.g., $1 + Y_{0,j,k}$) are understood as 1.

Then, $Y_{i,j,k+2(r+s+2)} = Y_{i,j,k}$ for all i, j, k .

- **Observation (Max Glick and others, ca. 2015?)**: This is equivalent to periodicity of birational rowmotion ($R^{p+q} = 1$) for $[p] \times [q]$, where $p = r + 1$ and $q = s + 1$, when the ring is commutative. Explicitly,

$$Y_{i,j,i+j-2k} = (R^k f)(i, j + 1) / (R^k f)(i + 1, j).$$

(Fine points omitted.)

- **Zamolodchikov periodicity conjecture in type AA** (proved by A. Yu. Volkov, [arXiv:hep-th/0606094v1](https://arxiv.org/abs/hep-th/0606094v1)): Let r and s be positive integers. Let $Y_{i,j,k}$ be elements of a commutative ring for $i \in [r]$ and $j \in [s]$ and $k \in \mathbb{Z}$. Assume that

$$Y_{i,j,k+1} Y_{i,j,k-1} = \frac{(1 + Y_{i+1,j,k})(1 + Y_{i-1,j,k})}{(1 + 1/Y_{i,j+1,k})(1 + 1/Y_{i,j-1,k})}$$

for all i, j, k , where sums involving “off-grid” points (e.g., $1 + Y_{0,j,k}$) are understood as 1.

Then, $Y_{i,j,k+2(r+s+2)} = Y_{i,j,k}$ for all i, j, k .

- **Observation (Max Glick and others, ca. 2015?)**: This is equivalent to periodicity of birational rowmotion ($R^{p+q} = 1$) for $[p] \times [q]$, where $p = r + 1$ and $q = s + 1$, when the ring is commutative.
- **Disappointment**: Zamolodchikov periodicity does not generalize to noncommutative rings (no matter how we order the five factors).

- A recent preprint by Joseph Johnson and Ricky Ini Liu (*Birational rowmotion and the octahedron recurrence*, [arXiv:2204.04255](https://arxiv.org/abs/2204.04255)) reproves the “order $p + q$ ” theorem for commutative \mathbb{K} in a simpler way (besides doing a number of other interesting things).

- A recent preprint by Joseph Johnson and Ricky Ini Liu (*Birational rowmotion and the octahedron recurrence*, [arXiv:2204.04255](https://arxiv.org/abs/2204.04255)) reproves the “order $p + q$ ” theorem for commutative \mathbb{K} in a simpler way (besides doing a number of other interesting things).
- The main idea of their proof is to reduce birational rowmotion to the octahedron recurrence, and prove the latter is periodic using lattice paths and LGV.

- A recent preprint by Joseph Johnson and Ricky Ini Liu (*Birational rowmotion and the octahedron recurrence*, [arXiv:2204.04255](https://arxiv.org/abs/2204.04255)) reproves the “order $p + q$ ” theorem for commutative \mathbb{K} in a simpler way (besides doing a number of other interesting things).
- The main idea of their proof is to reduce birational rowmotion to the octahedron recurrence, and prove the latter is periodic using lattice paths and LGV.
- We don't know if the octahedron recurrence is well-behaved for noncommutative \mathbb{K} (too many options to check), but LGV certainly is not available.

- A recent preprint by Joseph Johnson and Ricky Ini Liu (*Birational rowmotion and the octahedron recurrence*, [arXiv:2204.04255](https://arxiv.org/abs/2204.04255)) reproves the “order $p + q$ ” theorem for commutative \mathbb{K} in a simpler way (besides doing a number of other interesting things).
- The main idea of their proof is to reduce birational rowmotion to the octahedron recurrence, and prove the latter is periodic using lattice paths and LGV.
- We don't know if the octahedron recurrence is well-behaved for noncommutative \mathbb{K} (too many options to check), but LGV certainly is not available.
- Lemma 4.1 in the Johnson-Liu preprint generalizes our conversion lemma in the commutative case from single paths to k -tuples of nonintersecting paths. We don't know how this could be done in the noncommutative case; it is unclear in what order to multiply labels from different paths.

Proposition (2022, G & Roby):

Let P be any finite poset. Let $f \in \mathbb{K}^{\hat{P}}$. Then,

$$f(1) \cdot \sum_{\substack{u \in \hat{P}; \\ u > 0}} \overline{(Rf)(u)} \cdot f(0) = \sum_{\substack{u \in \hat{P}; \\ u < 1}} f(u),$$

assuming that the inverses $\overline{(Rf)(u)}$ are well-defined.

Proposition (2022, G & Roby):

Let P be any finite poset. Let $f \in \mathbb{K}^{\widehat{P}}$. Then,

$$f(1) \cdot \sum_{\substack{u \in \widehat{P}; \\ u > 0}} \overline{(Rf)(u)} \cdot f(0) = \sum_{\substack{u \in \widehat{P}; \\ u < 1}} f(u),$$

assuming that the inverses $\overline{(Rf)(u)}$ are well-defined.

Corollary (2022+, G & Roby):

Let P be any finite poset. Let $f \in \mathbb{K}^{\widehat{P}}$ with $f(0) = f(1) = 1$. Then, the quantity

$$\sum_{\substack{u, v \in \widehat{P}; \\ u < v}} f(u) \cdot \overline{f(v)}$$

is unchanged under birational rowmotion (i.e., when we replace f by Rf).